

## RESEARCH ARTICLE

# The interactive effects of temperature and food consumption on growth of larval Arctic cod (*Boreogadus saida*): A bioenergetic model

Carmen L. David<sup>1,2,\*</sup>, Rubao Ji<sup>2</sup>, Caroline Bouchard<sup>3,4</sup>, Haakon Hop<sup>5</sup>, and Jeffrey A. Hutchings<sup>1,6,7</sup>

Understanding larval growth, mediated by the interaction of early life traits and environmental conditions, is crucial to elucidate population dynamics. We used a bioenergetic model as an integrative tool to simulate the growth of Arctic cod (*Boreogadus saida*) larvae and to test the sensitivity of modeled growth to temperature and food quantity and quality. The growth was computed as the energy gained through food consumption minus the energy lost through respiration and other metabolic processes. We extended a previously published bioenergetic model to cover the full range of larval length and used a simplified feeding module. This simplification allowed us to build a predictive tool that can be applied to larval Arctic cod at a large spatial scale. Our model suggested that with subzero temperatures in the High Arctic, larvae need to increase food consumption in order to reach the observed length-at-age in late summer. The modeled growth agreed well with the field observations in the High Arctic but was 2–3 times higher than the laboratory-derived growth rate, probably due to differences in food type and selective mortality. Our study reveals important knowledge gaps in our understanding of larval cod growth in the High Arctic, including the lack of empirical estimations of daily ration and respiration for larvae under the natural habitat temperatures.

**Keywords:** Polar cod, Bioenergetic model, Polar fish, Feeding energetics, Temperature limitation

## 1. Introduction

Understanding the factors determining Arctic cod (*Boreogadus saida*) recruitment is a key element in predicting the response of the species to climate change in different parts of its circumpolar distribution. The Arctic cod is integral to Arctic ecosystems because of its high biomass and its importance as a food source for seabirds, fish, and marine mammals, especially in the High Arctic (Mecklenburg and Steinke, 2015; Mueter et al., 2020). The life cycle of Arctic cod is intimately connected to sea ice, suggesting a particular vulnerability of this species to changes in sea

ice, especially during its early life stages (Graham and Hop, 1995; Bouchard et al., 2017). Larval growth rate could directly influence survival probability during early life and, thus, recruitment to the adult population. Temperature and food consumption are the most important extrinsic factors of Arctic cod larval growth (Thanassekos et al., 2012; Koenker et al., 2018), but determining their relative importance and their interactive effects remains a great challenge, especially for wild larvae.

The reproductive strategy of Arctic cod includes a very long hatching season, extending from January to July (Bouchard and Fortier, 2008, 2011). Larvae from the same region and year but with different hatch dates can experience very different environmental conditions during their development, resulting in different growth and survival rates (Bouchard and Fortier, 2008, 2011). In region-years with environmental conditions allowing for the survival of winter hatchers, favored by increased food availability and lower predation, these individuals have an important advantage over the summer hatchers (Bouchard et al., 2017; LeBlanc et al., 2020). Despite slower growth during the first weeks of life resulting from low temperature, a longer growth season maximizes their prewinter weight and lipid reserves, important assets to surviving the first winter (Sogard, 1997; Copeman et al., 2020).

<sup>1</sup> Department of Biology, Dalhousie University, Halifax, NS, Canada

<sup>2</sup> Department of Biology, Woods Hole Oceanographic Institution, Falmouth, MA, USA

<sup>3</sup> Greenland Climate Research Centre, Greenland Institute of Natural Resources, Nuuk, Greenland

<sup>4</sup> Département de Biologie, Université Laval, Québec, QC, Canada

<sup>5</sup> Norwegian Polar Institute, Fram Centre, Tromsø, Norway

<sup>6</sup> Institute of Marine Research, His, Norway

<sup>7</sup> Centre for Coastal Research, University of Agder, Kristiansand, Norway

\* Corresponding author:  
Email: [carmen.david@dal.ca](mailto:carmen.david@dal.ca)

Temperature and food availability in the natural habitat affect larval Arctic cod growth by regulating their bioenergetic balance between the food intake and metabolic losses. However, the interaction or combined effects of temperature and food on growth remain unclear. Studies on larval fish at low temperatures have found that growth rates documented in the laboratory can be higher than those observed in the wild (see examples in Clarke and North, 1991; Icelandic Atlantic cod *Gadus morhua* in Björnsson, 1999). A key difference between laboratory and natural settings could be the food environment in terms of both quantity and quality. If optimal food conditions are not met by larval Arctic cod in the wild, then answering the question of whether temperature imposes an upper limit on the growth from the spatial distribution data alone is difficult (Clarke and North, 1991; Eriksen et al., 2020). Laboratory measurements of metabolic rates, such as respiration, heart rate, and feeding, suggest that larval Arctic cod exhibits a broader thermal tolerance than what is realized in their natural habitat (Drost et al., 2016; Kent et al., 2016). Therefore, to consider temperature as the determining factor of their natural habitat range is too simplistic. The challenge resides in considering the limitations imposed by both temperature and food on metabolic rates in order to be able to model larval growth at comparable levels to what is observed in the wild.

For larval Arctic cod, the relative sensitivity of growth to temperature and food consumption has not been directly compared over the full range of values observed in the field. A modeling study by Thanassekos and Fortier (2012), which used a bioenergetic model to simulate larval growth at different temperatures and food levels, predicted growth rates within the observed range in the field, but only for larvae smaller than 12 mm. Recent experimental work on early larval development in relation to temperature has revealed a growth optimum of 3°C–6°C (Koenker et al., 2018), surprisingly higher than estimated optimal temperatures derived from field observation data (Bouchard and Fortier, 2011). While the higher temperature optimum for growth might hold true for larger larvae (i.e.,  $\geq 12$  mm) and juveniles, previous studies reported that temperatures above 3.5°C result in poor hatching success and larval deformities (Sakurai et al., 1998; Kent et al., 2016). Moreover, direct comparisons between any kind of model predictions of growth patterns and observed growth of larvae in natural or seminatural environments are lacking for larval Arctic cod.

In the present study, we adapt and expand the bioenergetic model originally developed by Thanassekos and Fortier (2012) to cover the full-size range of larval Arctic cod and to test its applicability over different High Arctic regions. The model is used as an integrative tool for investigating the effects of temperature and food consumption on Arctic cod larval growth in order to understand the differences between experimental work and field observations. We hypothesize that (1) variations of 1°C–2°C at the lower temperature range can lead to large differences in size-at-age of larval Arctic cod by the end of their first summer and that (2) both food quantity, in terms of fraction of maximum daily ration consumed, and food quality, in terms of

energetic content of prey, are equally important in driving the observed variability of growth rates. This study will help us understand how the Arctic cod larval growth is affected by the interactions of early life history traits (e.g., hatching timing and size-at-hatch) and environmental conditions (e.g., temperature and food availability), which represents an important step in predicting how this Arctic species will respond to ongoing environmental changes.

## 2. Materials and methods

### 2.1. Sampling and aging of wild larval Arctic cod

Previously published length-at-age data of age-0 Arctic cod were estimated from otolith measurements of pelagic larvae and juveniles sampled in late summer and early fall between 2003 and 2015 in four regions of the Arctic Ocean (Bouchard and Fortier, 2011, 2020), for a total of 14 region–years. Fish were sampled from August 14 to October 8, standard length (SL) of fresh individuals measured, and otoliths extracted. A total of 1,092 fish were aged by otolith analysis (**Table 1**).

The sampling was conducted in three regions of the North American Arctic (Baffin Bay, Beaufort Sea, and Kitikmeot region) and one region of the Siberian Arctic (Laptev and East Siberian Sea, hereafter referred to as Laptev Sea). All sampled regions are located north of the Arctic Circle and ice-covered most of the year, with some spatiotemporal differences in timing of sea-ice retreat, ranging from May to September. Daily sea surface temperature (SST) in these regions ranges from near freezing –1.8°C immediately under ice cover to about a maximum 5°C in summer (**Figures 1** and S1–S4).

Daily SST fields were downloaded from the Estimating the Circulation and Climate of the Ocean Consortium (<https://www.ecco-group.org/home.cgi>; ECCO2 cube92 SST data set; Menemenlis et al., 2008) in each region–year, covering the estimated months of hatching and larval growth (from February 1 until September 15). The four regions were delimited geographically to include all Arctic cod sampling stations, as described in Bouchard and Fortier (2011) and LeBlanc et al. (2020).

### 2.2. Bioenergetic model

Bioenergetic models compute growth rate as a balance between food consumption (energy gain) and metabolic losses (respiration, excretion, and other losses through activity), both of which are affected by fish size and temperature (Brett, 1979; Railsback and Rose, 1999). These models can be used to synthesize metabolic processes and infer ecological relationships between fish and their environment (Ricker, 1979; Kitchell and Breck, 1980; Roy et al., 2004).

Here, we adapted a bioenergetic model (BEM hereafter) from Thanassekos and Fortier (2012; TF2012 model hereafter) to simulate Arctic cod larval growth over time and to quantify the effect of temperature and food consumption on larval growth. The model assumes that (1) daily consumption is a constant proportion of maximum consumption, (2) egestion and excretion are constant fractions of consumption, (3) energy content of fish and prey are constant, and (4) the ratio between active metabolic rate and basal standard metabolic rate is fixed. A major difference

**Table 1.** Arctic cod data used for parameter fitting and model validation. DOI: <https://doi.org/10.1525/elementa.2021.00045.t1>

Region	Year	Sampling Date	Number of Fish Aged	Mean $\pm$ SD SST <sup>a</sup>	Reference
Baffin Bay	2005	August 14–September 19	71	$-1.40 \pm 0.34$	Bouchard and Fortier (2011)
	2006	September 4–23	128	$-1.21 \pm 0.31$	Bouchard and Fortier (2011)
	2008	September 9–18	90	$-1.10 \pm 0.76$	Bouchard et al. (2017)
Beaufort Sea	2005	September 2–14	38	$0.69 \pm 1.36$	Bouchard and Fortier (2011)
	2010	August 15–24	197	$1.78 \pm 1.04$	Bouchard et al. (2017)
	2011	September 10–October 2	184	$1.21 \pm 0.83$	Bouchard et al. (2017)
	2014	August 18–September 9	80	$0.30 \pm 0.97$	Bouchard et al. (2017)
Kitikmeot	2005	August 27–30	32	$-1.33 \pm 0.16$	Bouchard et al. (2017)
	2006	September 25–27	12	$-1.23 \pm 0.28$	Bouchard et al. (2017)
	2011	October 6–8	23	$-0.98 \pm 0.39$	Bouchard et al. (2017)
	2015	August 15–September 23	29	$-1.39 \pm 0.29$	Bouchard et al. (2017)
Laptev Sea	2003	September 3–11	170	$-0.97 \pm 0.81$	Bouchard and Fortier (2008)
	2005	September 14–21	196	$-0.97 \pm 0.95$	Bouchard and Fortier (2008)
	2007	September 18–30	169	$-0.64 \pm 1.27$	Bouchard and Fortier (2011)

<sup>a</sup>Sea surface temperature (SST; Estimating the Circulation and Climate of the Ocean Consortium) shows average and standard deviation (SD) over the period April 15 to September 15 by region-year.

between our model and the TF2012 model is in the estimation of food consumption. First, a simplified formula, rather than a full forage module, was used in our estimation of consumption due to the lack of experimental data and associated parameter uncertainties. The simplified formula is similar to one used by Kitchell and Breck (1980) and Roy et al. (2004). The simplified consumption estimation in our model does not require input from field observations of prey data, which are not available for the entire growth season in large parts of the Arctic. Second, we revised the fraction of nauplii in the diet used in the TF2012 model, which was estimated for larvae <12 mm in length, and extended the formulation to the full age-0 length range (4–50 mm). To ensure the applicability of our model at the pan-Arctic scale, we validated our results with length-at-age estimations from otolith measurements from four High Arctic regions covering multiple sampling years (Table 1). The model predictions remain restricted to the High Arctic until further validation can be made with data on larval Arctic cod from more southern and warmer geographic areas.

Throughout this article, the length of fish refers to the fresh standard length (SL, mm) unless specifically stated otherwise. The conversion of wet weight ( $W$ ) to SL in the model is based on the regression  $W_{(g)} = 0.0055 SL_{(cm)}^{3.19}$ , provided by Geoffroy et al. (2016).

### 2.2.1. Growth rate

The weight-specific growth rate ( $G$ ,  $g\ g^{-1}\ d^{-1}$ ), here defined as daily weight change per unit of fish weight, can be

calculated based on the following energy-budget equation:

$$G = C \cdot A - R \cdot (f_{act} + SDA), \quad (1)$$

where  $C$  ( $g\ g^{-1}\ d^{-1}$ ) is the weight-specific consumption rate and  $A$  is the assimilation efficiency, representing the proportion of consumption remaining after egestion and excretion (Hop et al., 1997).  $R$  ( $g\ g^{-1}\ d^{-1}$ ) is the weight-specific rate of respiration and basal standard metabolism,  $f_{act}$  is a proportional increase in respiration due to activity, and  $SDA$  is the specific dynamic action. These terms are explained in detail in the sections below.

### 2.2.2. Consumption

The weight-specific consumption rate ( $C$ ,  $g\ g^{-1}\ d^{-1}$ ) is given by the formula:


$$C = C_{max} \cdot P, \quad (2)$$

where  $C_{max}$  is the maximum weight-specific consumption rate at weight ( $W$ ) and  $P$  is defined as a proportion (ranging from 0 to 1) of  $C_{max}$ . The maximum weight-specific consumption rate at weight ( $C_{max}$ ,  $g\ g^{-1}\ d^{-1}$ ) is defined as:

$$C_{max} = a_c \cdot W^{b_c} \cdot f(T) \cdot \left( \frac{CD_{prey}}{CD_{fish}} \right), \quad (3)$$

where  $a_c = 0.0337$  and  $b_c = -0.2862$  were estimated from the maximum daily ration formula:

$$\log_{10} MDR = 1.5272 + 0.7138 \cdot \log_{10} W, \quad (4)$$



**Figure 1. Sea surface temperature (SST) in the High Arctic in May 2020.** White polygons indicate the regions of Arctic cod samples used in this study: (a) Baffin Bay, (b) Beaufort Sea, (c) Kitikmeot, and (d) Laptev Sea (**Table 1**). SST data were downloaded from the National Oceanic and Atmospheric Administration (NOAA) data server ERDDAP, Estimating the Circulation and Climate of the Ocean (ECCO) Consortium, ECCO2 cube92 SST data set (Menemenlis et al., 2008). DOI: <https://doi.org/10.1525/elementa.2021.00045.f1>

experimentally determined for a size range of fish from juvenile to adult (weighing 3.99–59.19 g) at 0°C by Hop et al. (1997). The initial formula  $\log_{10} \text{MDR}$  ( $\text{mg}_{\text{prey}} \text{d}^{-1}$ ) was written in its exponential form, converted to grams and divided by fish weight ( $W$ ) to obtain the formulation  $\text{MDR}$  ( $\text{g}_{\text{prey}} \text{g}_{\text{fish}}^{-1} \text{d}^{-1}$ ) =  $a_c \cdot W^{b_c}$ , here corresponding to consumption as a function of fish weight.

The temperature function for the consumption  $f(T)$  was estimated by Thanassekos and Fortier (2012) from gut content volume ( $V$ ) of field-captured Arctic cod larvae. The functional response was fitted as the regression of  $V/V_{\text{max}}$  to temperature and standardized to 1 (where  $V_{\text{max}}$  is the maximum prey volume at larval length at 1-mm intervals), resulting in:

$$f(T) = \begin{cases} 0.943 + 0.155 \cdot T - 0.103 \cdot T^2, & T < 0.7^\circ\text{C} \\ 1, & T \geq 0.7^\circ\text{C} \end{cases} \quad (5)$$

The ratio of calorific density of prey ( $CD_{\text{prey}}$ ) to calorific density of larvae ( $CD_{\text{fish}}$ ) was added to express the efficiency of energy conversion from prey to fish tissue (Thanassekos and Fortier, 2012). By adding this ratio, the consumption becomes an expression of the proportional daily increase in fish weight ( $\text{g}_{\text{fish}} \text{g}_{\text{fish}}^{-1} \text{d}^{-1}$ ), eliminating the energy units. The calorific density of prey changes as larger larva selectively feed on larger prey, gradually

switching from a mixed diet of copepod nauplii ( $CD_n$ ) and eggs ( $CD_e$ ) to copepod nauplii alone:


$$CD_{\text{prey}} = \text{npl} \cdot CD_n + (1 - \text{npl}) \cdot CD_e. \quad (6)$$

The fraction of nauplii (npl) in the diet was estimated from gut content of larvae <12 mm by Thanassekos and Fortier (2012; **Table 2; Figure 2**). In addition to nauplii, larvae >15 mm gradually include copepodites in their diet; however, nauplii remain the dominant items in the diet of larvae <25 mm (Bouchard and Fortier, 2020). As both groups have similar calorific density, we did not specifically include the copepodites but assume that they are part of the diet as larvae increase in size and improve their swimming ability. Gradual increase in swimming speed with the size of both larvae and a larger prey would result in little difference in the outcome of any encounter as long as a size ratio of 1/10 is maintained (Fiksen and MacKenzie, 2002). Hence, the balance between energetic expenses of foraging, including failed attempts on larger prey, and energetic gain on captured prey would presumably be maintained.

The parameter  $P$  in Equation 2 was defined by Roy et al. (2004) as the proportion of  $C_{\text{max}}$  possible for a larva at a given temperature and prey availability ( $PA$ ,  $\text{g}_{\text{prey}} \text{g}_{\text{fish}}^{-1} \text{d}^{-1}$ ), where  $P = PA/C_{\text{max}}$ . When  $PA < C_{\text{max}}$ , consumption  $C$  becomes a fraction of  $C_{\text{max}}$ . Increased  $PA$  leads to higher

**Table 2.** Values of parameters used in the bioenergetic model. DOI: <https://doi.org/10.1525/elementa.2021.00045.t2>

Parameter Description (Unit)	Symbol	Baseline Value	Reference
Proportion of food consumption	$P$	0.73	This study
Assimilation efficiency	$A$	0.8	Hop et al. (1997)
$C_{\max}$ allometric factor	$a_c$	0.0337	Hop et al. (1997)
$C_{\max}$ allometric exponent	$b_c$	-0.2862	Hop et al. (1997)
Calorific density of Arctic cod ( $\text{cal g}^{-1}$ )	$CD_f$	1,327	Hop et al. (1997)
Calorific density of nauplii ( $\text{cal g}^{-1}$ )	$CD_n$	1,716	Thanassekos and Fortier (2012)
Calorific density of eggs ( $\text{cal g}^{-1}$ )	$CD_e$	475	Comita et al. (1966); Mauchline (1998)
Fraction of nauplii in diet	npl	$1 - e^{-0.198 \cdot (SL - 0.34)}$	This study
Length at yolk exhaustion (mm)	$L_{yex}$	8.5	Michaud et al. (1996)
Oxycalorific coefficient ( $\text{cal g O}_2^{-1}$ )	$Q_{ox}$	3,234	Hop and Graham (1995)
$\text{VO}_2$ allometric factor	$a_v$	0.0355	Hop and Graham (1995); Høleton (1974)
$\text{VO}_2$ allometric exponent	$b_v$	-0.1699	Hop and Graham (1995); Høleton (1974)
Specific dynamic action	SDA	0.375	Hop and Graham (1995)

**Figure 2.** Variation in length-at-age of Arctic cod larvae with increasing fraction of copepod nauplii in the diet.

To simulate growth, the fraction of copepod nauplii in the diet was estimated to be either a fixed value (0.6, 0.8, and 1) or adaptive (eq.), as a function of larval standard length (SL) in millimeter. The estimation of this function (eq.) is shown in the inset panel with red line, for which  $f_{\text{nauplii}} = 1 - e^{-0.198 \cdot (SL - 0.34)}$ . This function was fitted using the regression  $f_{\text{nauplii}} = -0.0042 \text{ SL}^2 + 0.1055 \text{ SL} + 0.1961$ , provided by Thanassekos and Fortier (2012), based on field estimations of gut content of larva <12 mm length (lower left portion of the black line), with the assumption that larva >15 mm feed exclusively on copepod nauplii and other larger prey (Bouchard and Fortier, 2020), for which  $f_{\text{nauplii}} = 1$  (upper right portion of the black line). DOI: <https://doi.org/10.1525/elementa.2021.00045.f2>

consumption until  $P$  reaches 1, as further increases cannot be ingested due to physiological constraints. For yolk-sac larvae,  $P$  is equal to 1, as yolk reserves constitute

endogenous resources for feeding, hence maximum daily consumption is assumed. Because the conversion of caloric content from prey to fish biomass does not have

relevance for yolk-sac larvae, which rely on endogenous feeding, the term  $CD_{\text{prey}}/CD_{\text{fish}}$  was excluded from the  $C_{\text{max}}$  formulation (Equation 3) for the duration of the yolk sac. When Arctic cod larvae reach a length of  $>8.5$  mm, they are considered to have entirely lost their yolk reserves (Michaud et al., 1996) and to rely exclusively on exogenous feeding, with consumption rate computed as a fraction  $P$  of the maximum consumption  $C_{\text{max}}$ . We use this simplification in the BEM by replacing a complicated foraging module with a single parameter  $P$ . Treated as an unknown, in the absence of data on PA, the parameter  $P$  can be determined by fitting the model to field observations.

### 2.2.3. Respiration and metabolic losses

The weight-specific standard respiration rate ( $R$ ,  $\text{g g}^{-1} \text{d}^{-1}$ ) is modeled as a function of weight and temperature, adjusted by the ratio of the oxycaloric coefficient ( $Q_{\text{ox}}$ ) to calorific density of fish larva ( $CD_{\text{fish}}$ ):

$$R = a_v \cdot W^{b_v} \cdot f(T) \cdot \frac{Q_{\text{ox}}}{CD_{\text{fish}}}, \quad (7)$$

where parameters  $a_v = 0.0355$  and  $b_v = -0.1699$  were estimated by combining a weight-specific respiration equation with a temperature-dependent respiration equation, the latter becoming the temperature function for the respiration  $f(T)$  in Equation 7. For  $f(T)$  formulation, oxygen consumption rates, reported as  $VO_2$  ( $\text{gO}_2 \text{g}_{\text{fish}}^{-1} \text{h}^{-1}$ ), were shown to increase linearly with temperature:

$$VO_{2(10)} = 0.042 + 0.011 \cdot T. \quad (8)$$

The oxygen consumption rates were measured for 5-month acclimated fish at temperatures of  $0.4^\circ\text{C}$ – $2.7^\circ\text{C}$  and standardized to a 10-g fish for comparison (Hop and Graham, 1995), using the following formula:

$$VO_{2(10)} = VO_{2(W)} \cdot W/10^{1-B}, \quad (9)$$

in which the weight exponent  $B = 0.8301$  originates from the weight-specific equation estimated by Holeyton (1974) at a constant subzero temperature of  $-1.5^\circ\text{C}$  for Arctic cod in the weight range of 0.6–122.0 g:

$$\log_{10} VO_{2(W)} = 0.8301 \cdot \log_{10} W - 1.0485. \quad (10)$$

Solving Equation 9, in which  $VO_{2(10)}$  is replaced with  $f(T)$  expression as in Equation 8,  $VO_{2(W)}$  ( $\text{gO}_2 \text{g}^{-1} \text{h}^{-1}$ ) can be written as a function of both weight and temperature and becomes the  $R$  expression in Equation 7. To this formulation, the oxycaloric coefficient ( $Q_{\text{ox}}$ ,  $\text{cal g O}_2^{-1}$ ) was added to convert weight units of  $\text{O}_2$  into calories, which divided with the calorific density of fish larva ( $CD_{\text{fish}}$ ) will translate respiration into proportional daily loss in weight ( $\text{g}_{\text{fish}} \text{g}_{\text{fish}}^{-1} \text{d}^{-1}$ ), thus using the same units as consumption and solving the main energy-budget equation (Equation 1). The oxycaloric coefficient was estimated considering the respiration substrate for feeding Arctic cod with a diet mainly on *Calanus* copepods (Hop and Graham, 1995, and references therein).

Because the weight-specific respiration was estimated for resting fish, an additional increase is required to

reproduce an active metabolism. When larvae are foraging, the standard metabolism is elevated, estimated by a factor of 2.0 for larval Atlantic cod (Lough et al., 2005); assuming that Arctic cod larvae are foraging during the daytime (12 of 24 h), the standard respiration rate was increased by a factor of  $f_{\text{act}} = 1.5$ . The additional energy expended on all activities of the body incidental to the ingestion, digestion, absorption, and assimilation of a meal was quantified by the SDA and was expressed as daily proportional increase in respiration by Hop and Graham (1995), where  $SDA = 0.375$ .

### 2.4. Parameter fitting and model validation

To find the best value of the parameter  $P$ , expressing the proportion of the maximum daily food consumption, length-at-age data from otolith measurements from four Arctic regions were used. Each of the individual region-year data sets shown in **Table 1** (we used 10 of the 14, as the year 2005 was not used in parameter fitting but kept for a second independent validation) was randomly split in two (60%, 40%) for either fitting parameter  $P$  or for model validation, respectively. The choice of 60%–40% was made gradually to increase the statistical power for fitting  $P$  of some limited data sets, which had only few data points. Improving the fitting of the parameter  $P$  ultimately led to an increase in modeling efficiency (EF), as explained in the next paragraph. For each region-year, the model was initialized with the minimum larval length sampled in that respective region-year and run for the number of days to cover the growth of the oldest larvae (i.e., age in days of the oldest larvae minus the age of the youngest). The model used regionally averaged temperature (as showed in **Table 1**), and a range of  $P$  values from 0.5 to 0.9 (at 0.01 intervals) were tested.

For each value of  $P$  tested, an EF metric was calculated based on the formula  $EF = 1 - \sum |y_o - y_p|^2 / \sum |y_o - y_m|^2$  (Mayer and Butler, 1993), where  $y_o$  are the length-at-age values observed,  $y_p$  are the values predicted, and  $y_m$  is the mean of observation values. The  $P$  value with the highest EF was selected for the respective region-year. Then, using this value of  $P$ , the model was run again, and EF was calculated, using the second subset kept for model validation, representing 40% of the data in that respective region-year. Due to high uncertainty in the daily location of the sampled larvae, which in the High Arctic is on the order of a few hundred kilometers over the growth period (Eriksen et al., 2020; Vestfals et al., 2021), the choice was made to use regionally averaged temperature for model simulations. Moreover, different hatching cohorts can also experience different temperatures, such as early hatcher versus late summer hatcher. To override the growth differences arising from different hatching cohorts, the second choice made in this regard was to use an average temperature over the growth period of the dominant cohort, that is, the April hatchers (mean SST values between April 15 and September 15, shown in **Table 1**).

We acknowledge the potential errors associated with using an average temperature to simulate growth over few months. Therefore, we used an additional data set, which is the year 2005 in all four High Arctic regions (four

region–year data sets), to evaluate the model efficiency in predicting growth for different hatching cohorts using daily temperatures. For this validation, an average value of  $P$  was used per region and growth was initialized with four larval lengths, 4, 5, 6, and 7 mm, covering the hatching size variability of larval Arctic cod (Ponomarenko, 2000). The model was run using regionally averaged daily temperatures from the ECCO2 model (Menemenlis et al., 2008) and simulating larval growth with different daily hatching dates, starting from February 15. All simulations ended on September 15. EF was calculated for each region, using the predicted lengths within the same age interval as observations.

### 2.5. Sensitivity analysis

The model sensitivity analysis was conducted to assess the relative importance of parameters to the modeled growth rates. Here, we focus the analysis on the temperature and the parameters related to food quantity ( $P$ ) and quality (i.e.,  $CD_m$ ,  $CD_e$ ,  $CD_f$ ). Sensitivity analysis for the full list of parameters can be found in Supplemental Material (Texts S2 and S3). First, we quantified the sensitivity of growth rate to variation in model parameters through individual parameter perturbations (IPP; Bartell et al., 1986). Each parameter value was varied  $\pm 20\%$  of the baseline value at a time while the other parameters were held constant (Table 2). For temperature to be representative of the High Arctic regions, a reference temperature of  $0^\circ\text{C}$  was chosen which then was varied  $\pm 1^\circ\text{C}$ . Variation in larval length after 30 days of growth, initialized with a size at hatch of 6 mm, was compared to a reference larval length simulated using the nominal values of all parameters. The relative change in modeled length ( $L_m$ ) from the reference ( $L_{\text{ref}}$ ) was estimated as the ratio  $\Delta L = (L_m - L_{\text{ref}})/L_{\text{ref}}$ . Second, we applied a global sensitivity analysis in which all parameters were varied simultaneously, each randomly selected from a normal distribution. We implemented this method by using Latin Hypercube Sampling (van Griensven et al., 2006) and determined the relative contribution of each parameter variation to prediction error (Bartell et al., 1986). Normal distributions were assigned to the parameters included in the IPP analysis by using the nominal values as the mean and their  $\pm 20\%$  variation as the minimum and maximum values (Table 2). Values were drawn randomly from the distributions to create a set of parameter values in each model simulation. Each parameter distribution was sampled 300 times, resulting in a data set  $300 \times 6$  (five parameters and modeled larval length). The data set was analyzed statistically by calculating Pearson correlations between simulated larval length and each parameter. The relative contribution of each parameter to prediction error was estimated by multiple linear regression and a bootstrapping method. We repeated the analysis for two sizes of larva (SL of 6 and 12 mm).

### 2.6. Variations in food quantity and quality

Different simulation scenarios were cross-tested in which (1) food quantity and (2) food quality were modified. The change in food quantity was simulated by using a fraction of maximum daily food consumption of  $P = 0.6$  which

then was reduced to  $P = 0.3$ . The alteration in food quality was implemented in the model by using either a diet based on a *Calanus nauplii* caloric content of  $1716 \text{ cal g}^{-1}$  (Table 2) or a diet based on *Artemia nauplii* of  $1,295 \text{ cal g}^{-1}$  (equivalent to  $20.9 \text{ kJ g}^{-1}$  according to Vanhaecke et al. (1983), considering  $1 \text{ cal} = 4.1868 \text{ J}$  and a water content of  $74\%$ ). In all scenarios, the BEM was initialized with a length-at-hatch of 6 mm and used a constant temperature of  $0^\circ\text{C}$  during growth. The resultant modeled length-at-age was compared among different scenarios, informed by both field measurements and laboratory experiments.

## 3. Results

### 3.1. Parameter $P$ variability and model validation

Within the same region, the best values estimated for the parameter  $P$  were in good agreement across the years (Table 3). Higher average  $P$  per region of 0.85 and 0.82 were estimated for Baffin Bay and Kitikmeot region, the two regions with the coldest temperatures, followed in decreasing order by Laptev Sea with 0.70 and Beaufort Sea with 0.60. The implementation of these values of  $P$  in the validation runs resulted in an EF between 0.6 and 0.97, with the lowest values associated with the region–years having the lower number of observations, irrespective of region (Table 3). This association was also noticeable in the visual comparison of observations versus predicted values of length-at-age (Text S1, Figures S5–S14).

Modeled length-at-age estimated using a fixed value of  $P$  per region and a constant temperature, averaged over the years, was well within the range of observations based on otolith measurement from four Arctic regions, when compared to individual regressions fitted per region and year (Figure 3).

When daily temperatures were used to simulate growth of larvae with different hatching dates, starting from February 1, the model responded well to the seasonal changes in temperature (Figure 4, showing only monthly cohorts), with EF falling between 0.44 and 0.94, except for the Beaufort Sea. The model performed poorly in Beaufort Sea, where the growth patterns seemed more distinct between different monthly cohorts (Figure 4). The modeled lengths-at-age in Beaufort Sea were underestimated for the early hatchers with longer growth period and overestimated for the late hatchers. For the other regions, higher EF per each region in the 2005 validation simulations was correlated to different lengths-at-hatch, namely, 4 mm for Baffin Bay (EF = 0.80), 6 mm for Kitikmeot region (EF = 0.94), and 7 mm for Laptev Sea (EF = 0.76; Table S1).


### 3.2. Sensitivity of growth to model parameters

Modeled growth was sensitive to the group of parameters involved in food consumption, when these were individually varied with  $\pm 20\%$  of their nominal values (Figures 5 and S15, Text S2, Table S2). The fraction  $P$  of  $C_{\text{max}}$  and  $CD_n$  resulted in a proportional change in length ( $\Delta L$ ) of  $\pm 0.08$  after 30 days of growth, while  $CD_f$  induced a similar but negative response in  $\Delta L$  of 0.08 and 0.05, respectively. The lower changes were induced by the temperature and

**Table 3.** Fitting the parameter for proportion of consumption ( $P$ ) and model validation. DOI: <https://doi.org/10.1525/elementa.2021.00045.t3>


Region	Year	Nf	$P$	Nv	EF
Baffin Bay	2006	76	0.85	52	0.611
Baffin Bay	2008	54	0.85	36	0.657
Beaufort Sea	2010	118	0.60	79	0.794
Beaufort Sea	2011	110	0.61	74	0.730
Beaufort Sea	2014	48	0.60	32	0.681
Kitikmeot	2006	7	0.85	5	0.591
Kitikmeot	2011	272	0.85	182	0.971
Kitikmeot	2015	355	0.76	237	0.838
Laptev Sea	2003	102	0.69	68	0.791
Laptev Sea	2007	101	0.70	68	0.809

Nf = number of observations used for parameter fitting;  $P$  = proportion of maximum daily food consumption; Nv = number of observations used for model validation; EF = modeling efficiency.



**Figure 3.** Fraction of the maximum daily ration  $P$ . Estimation of the parameter  $P$  was done by region–year based on data from otolith measurements of larval Arctic cod (see Bouchard and Fortier, 2011, for details). Regressions based on otolith measurement data are shown by continuous line and bear the name of the respective region in the legend of each panel. The model output using a  $P$  value of 0.60, 0.71, 0.82, and 0.85, indicated by  $P$  and solid line, was initialized with a length at hatch ( $L_{ini}$ ) estimated from otoliths regressions, averaged over the years indicated in the legend, and using an average field surface temperature ( $T$ ) as shown in each panel (Table 1). Data points represent the field observations of length-at-age in the respective region–year. DOI: <https://doi.org/10.1525/elementa.2021.00045.f3>






**Figure 4. Length-at-age for multiple hatch-date cohorts in 2005.** Growth was simulated with the bioenergetics model. Simulations were initialized with a length at hatch of 6 mm and different hatch dates (shown in the legend) and used daily SST starting from the hatch date until September 15, 2005, averaged per region. Data points (“observations” in the legend) represent length-at-age of fish caught in 2005 in the four regions and aged by otoliths measurements (Table 1). Number of fish aged per region in 2005 is shown in parentheses. For each region, the value of  $P$  used in the model is indicated and the modeling efficiency estimated. DOI: <https://doi.org/10.1525/elementa.2021.00045.f4>

$CD_e$  with a proportional  $\Delta L$  from the reference length  $<0.01$  when the parameters values were increased (Figure 5B). When the values were lowered, however, the most sensitive parameter was temperature, resulting in negative  $\Delta L$  of 0.13 of the reference length (Figure 5B).

When a global sensitivity analysis was applied to test the combined effect of the temperature and food-related parameters, the same group of parameters sensitive to the IPP induced the largest changes in model output (Figures 6A–E and S16–S18, Text S3). When both temperature and food-related parameters were varied, with values randomly drawn from a normal distribution, modeled larval length by the BEM showed a significantly negative trend across the temperature range tested (Figure 6A, Loess regression line). Among food-related parameters,  $P$  and  $CD_n$  showed a significantly positive linear effect on modeled length (Figure 6B and D). Each parameter contributed between 10% and 39% of the overall response variance ( $R^2 = 77\%$ , metrics were normalized to sum

100%), with  $CD_n$  and  $P$  having the largest contribution (Figure 6F).

When metabolic rates of larvae were simulated over the range of temperatures found in the High Arctic and tested in experiments (up to  $9^\circ\text{C}$ ; Koenker et al., 2018), growth rates were more sensitive to and increased sharply across the negative temperatures, followed by the gradually decreasing trend due to increase in respiration at higher temperatures (Figure 7A). At negative temperatures, growth rates were less sensitive to variations in the proportion  $P$  of the maximum daily ration ( $P = 0.73$  and  $\pm 20\%$  range for values of  $P = 0.58$ – $0.87$ ). Modeled growth fell within the range of laboratory estimates at higher temperatures, considering variations in the parameter  $P$ . At temperatures  $<3^\circ\text{C}$ , our modeled growth rates were 2–3 times higher than laboratory estimates. To maintain a positive growth rate, a larva of 12-mm length requires a minimum growth value of  $P = 0.12$ – $0.15$  over the negative temperatures (Figure 7B). For the positive temperatures, the minimum value of  $P$  required for positive



**Figure 5. Sensitivity analysis by individual parameter perturbation (IPP).** (A) Length-at-age from different model outputs following the IPP analysis, in which parameter values were increased by 20% their nominal values and the temperature used was 1°C. In all simulations, larval growth was initialized with a size-at-hatch of 6 mm and used a temperature of 0°C. *Ref* represents larval length over a 30-day period of growth using the nominal values of all parameters. (B) The ratio of change in larval length from the reference length (*Ref*) after 30 days of growth, when each individual parameter was varied  $\pm 20\%$  at the time, and low/high temperatures were set at  $-1^\circ\text{C}$  and  $1^\circ\text{C}$ . The parameters are ordered from higher to lower sensitivity (left to right). DOI: <https://doi.org/10.1525/elementa.2021.00045.f5>

growth increased linearly with temperature from 0.12 at  $0^\circ\text{C}$  to 0.45 at  $10^\circ\text{C}$ .

### 3.3. Sensitivity of modeled growth to food quantity and quality

Modeled length by the BEM for a 30-day-old larva was in the upper range of field estimates and 1.5–4.5 mm higher than the growth observed in laboratory experiments (Figure 8). When yolk reserves were present, which lasted about 6–13 days at  $0^\circ\text{C}$ , depending on the length-at-hatch, the model overestimated growth. This output is in comparison with previous observations of yolk-sac duration in Arctic cod larvae from hatching until complete absorption, which lasts about 20 days at  $1^\circ\text{C}$  (Aronovich, 1975; Graham and Hop, 1995). In the post-yolk period, variations in modeled growth and resultant length-at-age were related to both changes in food quantity (through

variations of  $P$ ) and quality (through variations in caloric density of prey). When a 25% lower caloric diet based on *Artemia* nauplii ( $1,295 \text{ cal g}^{-1}$ ; Vanhaecke et al., 1983) was simulated instead of a richer *Calanus* nauplii diet ( $1,716 \text{ cal g}^{-1}$ ; Thanassekos and Fortier, 2012, and references therein), which is implicitly used in the BEM, a 10% decrease in length-at-age resulted in the model output. This value remains higher than laboratory observations on larval Arctic cod growth. Reducing the daily ratio of food consumption in the model by 50% resulted in similar growth rates as laboratory estimates.


## 4. Discussion

### 4.1. The effect of food consumption on growth

The sensitivity analysis of the model parameters indicated that those describing the feeding process are responsible for the largest changes in model predictions, a finding consistent with other bioenergetic models (Lough et al., 2005; Kristiansen et al., 2007). This finding is due, first, to order-of-magnitude higher food consumption rates compared to respiration rates and metabolic losses. Higher sensitivity in our modeled growth related primarily to variations in the proportion  $P$  of the maximum daily ration ( $C_{\text{max}}$ ). Second, the energy of prey ingested by the larvae is not only a matter of quantity but also of food quality, reflecting the energetic conversion efficiency from prey to fish tissue (Støttrup, 2000).

In the BEM formulation, the energetic conversion efficiency was included as a ratio of the caloric content of *Calanus* nauplii/copepodites as prey to the caloric content of fish (Thanassekos and Fortier, 2012). At a lower energetic content of food, this ratio is reduced, hence less energy is gained. A diet of enriched *Artemia* nauplii, commonly used in laboratory experiments as food source for fish larvae (Graham and Hop, 1995; Koenker et al., 2018), has about 25% lower caloric content than a diet comprised of *Calanus* nauplii and copepodites (Vanhaecke et al., 1983; Sorgeloos et al., 2001), the dominant food source for larval Arctic cod in the High Arctic (Kohlbach et al., 2017; Bouchard and Fortier, 2020). Simulating this reduction in the caloric intake in the model resulted in a 10% decrease in the length of the early-feeding larvae at 30 days posthatching. Variation in growth related to food type has also been observed in yearling (age 1+) Arctic cod. Fish fed *Calanus* copepods had significantly higher growth rate compared to other food sources such as *Themisto* amphipods or capelin filets (Hop et al., 1997). Such diet differences may have important consequences for larval development. The consequences of poor nutrition during larval development may be obvious, for example, through lower growth and survival, deformities or malpigmentation (Shields et al., 1999; Payne and Rippingale, 2000), but in many cases, the consequences may be obscure, as in effects on temperature tolerance or limited growth during later life stages (Støttrup, 2000).

First-feeding Arctic cod larvae consume mainly copepod eggs and nauplii (80% of the stomach content) for the first days to weeks, switching gradually to a diet that includes copepodites with increased larval ability to




**Figure 6. Pearson correlation between larval length with temperature and food-related parameters.** Modeled larval length after 3 days of growth given variations in temperature and food-related parameters (A–E). Larval length was initialized with 12 mm. Linear regressions are shown in black and loess regression (locally weighted smoothing) in red. Random values for temperature were drawn by Latin Hypercube Sampling from a uniform distribution (min. =  $-1.8^{\circ}\text{C}$ , max. =  $1^{\circ}\text{C}$ ) and for the other parameters from normal distributions around their mean  $\pm 20\%$ . (F) Percentage of variance (% of  $R^2$ ) in a multiple regression model on larval length explained by variation in model parameters related to food quality and quantity and temperature with 95% confidence intervals. Model parameters, using variation in values drawn from a normal distribution by Latin Hypercube Sampling, were used as explanatory variable in the regression model, and larval length, simulated with the bioenergetics model, was used as response variable. Results were normalized to sum 100% ( $R^2 = 71.66\%$ ). DOI: <https://doi.org/10.1525/elementa.2021.00045.f6>

capture larger prey size (Walkusz et al., 2011; Bouchard and Fortier, 2020). A ratio of 1/10 was suggested between larva attack speed ( $10 \text{ SL s}^{-1}$  estimated for Atlantic cod) and prey escape speed ( $100 \text{ SL s}^{-1}$ ) in the calculation of capture success (Fiksen and MacKenzie, 2002). This ratio simulates a prey-size preference of  $<5\%$  of larval body length, in agreement with observations on Arctic cod diet by Bouchard and Fortier (2020) and on Atlantic cod diet by Munk (1997). As long as this ratio is maintained, swimming speed of both predator and prey would increase with size, resulting in little difference in the outcome of any encounter; hence, the balance between energetic spending on foraging and energetic gain on prey captured would presumably be maintained (Folkvord, 2005; Lough et al., 2005).

At high prey density, feeding success increases while the time spent foraging decreases, hence less energy spent

(Puvanendran and Brown, 1999; Monk et al., 2006). We assumed an average daily feeding ration over the growth season mediated through the parameter  $P$ , which was fitted to best simulate the observed growth of wild larvae in four High Arctic regions, which represent the survivors at the end of summer. Our model output reflected well field observations, even though it relied on a fixed  $P$  value per region-year; hence, it did not account for the spatiotemporal variations in prey density. In reality, suboptimal prey concentrations during the first feeding result in slower growth, which cannot be sustained for more than 2–3 weeks, resulting in higher mortality (Monk et al., 2006; Kristiansen et al., 2007; Koenker et al., 2018). Therefore, isolated records on positive growth are not necessarily indicative of longer term consequences or survival in a patchy prey environment (Laurel et al., 2018). While it seems intuitive, larval growth is not always correlated with



**Figure 7. Modeled metabolic rates and laboratory-derived growth rate with temperature.** (A) The effect of temperature on metabolic rates (consumption, *C*, and respiration, *R*) and growth (*GR*) estimated by the bioenergetics model (BEM). Simulations are shown for a larval fish of 12-mm length. Modeled growth rate, shown by a red line, used an average value of  $P = 0.73$ ; the uncertainty around the line indicates variations induced by changes in  $P$  value within the range of  $\pm 20\%$ . BEM uses an expression of temperature limitation provided by the experimental work of Hop et al. (1997), as used in the model presented by Thanassekos and Fortier (2012); see Equation 3 for *C* and Equation 7 for *R*. *Lab* uses a growth model regression from Koenker et al. (2018). (B) The minimum value of  $P$  required to maintain a positive growth rate over the range of temperatures shown by black line for a larval fish of 12-mm length. The linear regression equation for temperatures  $>0^\circ\text{C}$  is provided. The shade around the line indicates the minimum  $P$  value for the full length range of feeding larvae (8.5–27 mm). DOI: <https://doi.org/10.1525/elementa.2021.00045.f7>

prey density in the natural habitat (Michaud et al., 1996). Increasing prey density, on the other hand, can lead to better survival because larvae prioritize their predator avoidance while still maintaining growth (Fiksen and Jørgensen, 2011). Therefore, predicting survival, unlike growth, needs to consider the patchiness of prey and predators in the natural environment as well as larval behavior related to predator avoidance.


#### 4.2. The effect of temperature on growth

Environmental temperature is the most important variable affecting metabolic rates of fish larvae (Ikeda, 2016). In theory, larval growth increases with temperature as long as prey occur at sufficient density to offset the increased metabolic costs (Clarke, 2006). In our BEM, food consumption increases asymptotically with temperature, based on estimations by Thanassekos and Fortier (2012) on the temperature effect on stomach fullness in wild fish larvae, while respiration increases linearly, reaching a maximal modeled growth at an optimum temperature of  $0.7^\circ\text{C}$ . Laboratory estimates, on the other hand, indicate an optimal growth temperature of  $5^\circ\text{C}$  (Koenker et al., 2018). The large difference in optimum temperature is not only a result of model formulation but also represents an image of data used in model parametrization. Wild larvae sampling covered only a spatiotemporal snapshot of the High Arctic, while experimental work has its own limitations, such as temperature range, duration, or acclimation potential. These aspects should be kept in mind in model evaluation and comparison. When the same temperature range will be covered by field sampling and experimental work, then different model formulations can be better tested and compared.

Modeled growth by BEM in all four High Arctic regions was sensitive to variations in temperature as small as  $1^\circ\text{C}$ . As a result, the model predicted different growth curves for larvae with different hatch dates. This aspect is important to consider, as the model needs to capture accurately the seasonal increase in temperature from near freezing at the beginning of the hatching period to barely above zero when simulating growth of larvae in the High Arctic. However, field observations on wild larval growth were limited to High Arctic regions, while similar observations from warmer regions remain scarce. Laboratory experiments concur with the increase growth with temperature, but their key limitation resides in the difficulty of simulating the lower thermal range observed in the natural habitat, that is, the subzero temperatures (Graham and Hop, 1995; Koenker et al., 2018). Nevertheless, laboratory experiments emphasized the need to account for both species- and stage-specific differences in the thermal response of growth models (Koenker et al., 2018). Consequently, models are useful for bridging these thermal gaps between field and laboratory observations.

#### 4.3. Interacting effects of food consumption and temperature

While the relationship between respiration and temperature is relatively well known in fish, suggesting a positive and monotonic increase, close to exponential in shape (Clarke, 2006), estimations of food consumption with temperature remain scarce. This relationship is even more important to define, as variations in food consumption with temperature can largely dictate the optimum thermal window and growth patterns in fish larvae, as shown in our model. At higher than optimum temperatures for growth, exponential increase in respiration rates results in hyperventilation (Drost et al., 2014, 2016), while food uptake is increasingly used to cover the exponentially



**Figure 8. Length-at-age of posthatching larvae compared among bioenergetic model (BEM) simulations, field measurements and laboratory experiments.** The BEM projected growth (with an initial length-at-hatch of 6 mm), using a constant temperature of  $0^{\circ}\text{C}$  and a ratio of maximum daily food consumption  $P = 0.6$  in the upper panels (A and B) and  $P = 0.3$  in the lower panels (C and D). *FieldGR* uses a constant growth rate of  $0.20\text{ mm d}^{-1}$  based on otoliths measurements by Bouchard and Fortier (2011; range of  $0.18\text{--}0.21\text{ mm d}^{-1}$ ). *Field30* shows a boxplot of 30-day-old larval length, estimated from regional length-at-age regressions. *Lab1GR* uses a quadratic equation as a function of temperature from Koenker et al. (2018), which here at  $T = 0^{\circ}\text{C}$  growth rate becomes  $2.35\%$  body weight  $\text{d}^{-1}$ ; *Lab2GR* uses a growth rate of  $0.061\text{ mm d}^{-1}$ , estimated by Graham and Hop (1995) at  $1.5^{\circ}\text{C}$ . Left column (A and C) shows simulated growth by BEM using a diet based on *Calanus* copepod caloric content ( $1,716\text{ cal g}^{-1}$ ) and the right column (B and D) using a diet based on *Artemia* nauplii ( $1,295\text{ cal g}^{-1}$ ; Vanhaecke et al. (1983), which was the food source used in both laboratory experiments *Lab1GR* and *Lab2GR*. DOI: <https://doi.org/10.1525/elementa.2021.00045.f8>

higher metabolic demands, finally resulting in declining growth rates (Brett, 1979).

Surprisingly, observations on immature adult Arctic cod (age 1+ and 2+) indicated that growth rate did not vary significantly over the temperature range  $0^{\circ}\text{C}\text{--}8^{\circ}\text{C}$ , while respiration was strongly affected by temperature (Kunz et al., 2016). This finding implies that fish could have indeed compensated for increased respiration with higher food uptake. This assumption fits our results on the minimum proportion  $P$  of the maximum daily ration required to maintain positive growth. Nevertheless, this strategy for growth would then suggest that at least immature adult Arctic cod have the physiological potential to expand their distribution southward into warmer areas, given that enough food is available in the environment to compensate for increasing metabolism related to higher temperature. Although when the narrow range of thermal tolerance of Arctic cod juveniles overlapped with that of subpolar fish (e.g., saffron cod *Eleginus gracilis*), growth and lipid accumulation exceeded that of their southern

congeners only at water temperatures  $<2^{\circ}\text{C}$  (Laurel et al., 2016; Copeman et al., 2020), indicative of limiting factors of Arctic cod distribution other than temperature and food availability (probably food type and quality). This compensatory strategy for growth would further imply that at low temperature, Arctic cod require less food than at higher temperature or than southern competitor species to maintain similar growth rates, thus presenting an advantage in the High Arctic (Laurel et al., 2018).

Generally, fish larvae from cold ecosystems are believed to require less food than larvae from warm ecosystems (Pörtner et al., 2005). This strategy does not concur with our modeled growth in four Arctic regions, which showed that larger daily ration (mediated through parameter  $P$ ) was needed in regions with lower temperature to reproduce the field-observed growth. A limitation in our models arises from the data used for parameter fitting and validation, which was restricted to the High Arctic region, north of the Arctic Circle, with a surface temperature range from  $-1.8^{\circ}\text{C}$  to  $5^{\circ}\text{C}$  at best (Figures 1 and

S1–S4). Furthermore, larvae are more thermally sensitive than juvenile and adult Arctic cod (Drost et al., 2016; Laurel et al., 2017) and could have a different strategy for optimal growth or for survival, upon which we can only speculate. Clear, however, is that these larvae have an adaptation potential that makes them thrive at near-freezing temperature.

Ultimately, the combined effects of increasing metabolic demands with temperature and compensation with food consumption are not only a matter of food quantity but of energetic gain. At low temperature, fish have lower metabolic demands, spend less energy foraging but have high energetic gain when a diet of lipid-rich Arctic prey is consumed (Kohlbach et al., 2016, 2017). Therefore, even when PA is limited, low temperatures could still yield the highest growth for Arctic cod in the High Arctic. Our model indicated that growth was less sensitive to variations in  $P$  at negative temperatures, implying that a relatively constant growth could be maintained despite spatiotemporal variations in PA, likely because at low temperatures, slow digestion limits food intake (Hop and Tonn, 1998). At high levels of available prey, maximum growth is normally expected to occur at higher temperatures (Lough et al., 2005). This dynamic response of metabolic processes to temperature agrees with laboratory and field observations on the growth of other fish species (Brett and Groves, 1979; Otterlei et al., 1999). For Arctic cod larvae, however, such growth dynamics remain speculative because of the lack of direct observations.

#### 4.4. Modeled growth

Modeled growth rates in larval Arctic cod resulted in similar rates to those observed in the wild and estimated from otolith microstructure (Bouchard and Fortier, 2011; Bouchard et al., 2017), except for the yolk-sac period when growth was overestimated by the model, predicting a shorter duration (Graham and Hop, 1995; Kent et al., 2016). Of interest is that the experimental work published to date on the growth of larval Arctic cod reported lower growth rates than those estimated by our model or observed in the wild at similar temperatures (Graham and Hop, 1995; Koenker et al., 2018; Bender et al., 2021). Our BEM might overestimate larval growth compared to laboratory settings for two reasons. First, food quality is better in nature (by 25% higher caloric content), with a variety of lipid-rich prey being available in the High Arctic (Kohlbach et al., 2016). Second, the model was adjusted using data from wild age-0 fish collected between August and October; that is, the survivors of the larval cohorts remaining after selective mortality likely removed the smaller-at-hatch and slower growing individuals from the population through predation and starvation (Peck and Hufnagl, 2012; Jørgensen et al., 2016). Selection for large size-at-hatch has been documented in larval Arctic cod (Pepin et al., 2014), while selection for fast growth is likely to occur among larvae hatching at the same time. Faster growth in wild larvae compared to reared larvae has been observed in other species, notably Atlantic cod (Finn et al., 2002; Folkvord, 2005).

To date, we are unaware of bioenergetic models of larval Arctic cod other than the one presented by Thanasekso and Fortier (2012), which derived daily ration estimates from immature adults in the absence of data on larvae. In the absence of accurate field prey data, model approximations can be informative despite their limitations, in the same way as experimental work that uses two fixed settings of prey density (low vs. high scenarios) over the growth period (Koenker et al., 2018). Given the variability in model validation studies and the uncertainty in model inputs (such as oxycaloric equivalents and the energy content of predators and their prey, among others), testing models at all empirically defensible thermal, size, and feeding levels, even in the laboratory, may not be possible. Our models, based on average daily values of  $P$  and regional temperatures, might have masked the potential for patchiness in prey concentration and temperature that could potentially boost Arctic cod consumption and growth or, on the contrary, lead to starvation.

Feeding behavior continues to be a difficult process to model and questions related to prey preference and foraging are particularly challenging (Fiksen and MacKenzie, 2002). Equations including individual foraging behavior (Kristiansen et al., 2007; Peck and Daewel, 2007) and depth-varying conditions (Fiksen and MacKenzie, 2002) or coupled biophysical simulations on match–mismatch feeding dynamics (Kristiansen et al., 2009) would perhaps account for small-scale variability in prey abundance and temperature. The increased accuracy in predicting growth provided by more sophisticated schemes, however, would come at higher computational cost. Moreover, field observations of prey data are required for such simulations, which are largely unavailable in the High Arctic, especially during the ice-covered period, therefore limiting the capacity of such a model to resolve, for example, the impact of hatch dates on the growth trajectory of different cohorts at a circum-Arctic scale. Alternatively, prey data could indeed be estimated from biochemical NPZD (nutrient, phytoplankton, zooplankton, and detritus) or ecosystem models, which come, however, with additional prediction errors that need to be considered. Thus, the key advantage in simplifying the feeding processes in a bioenergetic model, such as the BEM presented here, relies on the possibility to be further applied at larger spatial scales to test larval growth dynamics with limited input data.

#### 5. Conclusions

The bioenergetic model presented here provides a robust description of the general pattern in larval Arctic cod growth, synthesizing metabolic processes that are of both theoretical and practical importance to modelers and physiologists. With limited information on feeding behavior and underlying relationships between metabolic processes and temperature, BEM proved useful in estimating growth of larval Arctic cod, well within the range of field observations.

Modeling the variability of individual growth and its determining factors can considerably increase our understanding of Arctic cod recruitment, population dynamics, and response to environmental changes. Our model allowed us to test hypotheses with only limited information

at hand on prey availability or feeding dynamics and to identify crucial knowledge gaps, such as the need for estimating daily ration with larval weight and temperature and yolk-consumption dynamics. Further consideration should also be given to different types of food used in experimental work. Aquaculture feed typically is lower in quality than Arctic copepods and may lead to suboptimal growth even with high availability. Ultimately, using modeling tools to bridge gaps between laboratory and field studies can prove crucial in expanding our understanding of important species in Arctic seas.

### Data accessibility statement

The R scripts generated in this study for model formulation, data analysis, and figure plotting are publicly available on Github and can be accessed at [https://github.com/carmenldavid/ArcticCod\\_BioenergeticModel](https://github.com/carmenldavid/ArcticCod_BioenergeticModel).

### Supplemental files

The supplemental files for this article can be found as follows:

**Figure S1.** Sea surface temperature (SST) in Baffin Bay.  
**Figure S2.** Sea surface temperature (SST) in Beaufort Sea.

**Figure S3.** Sea surface temperature (SST) in Kitikmeot region.

**Figure S4.** Sea surface temperature (SST) in Laptev Sea.  
**Text S1.** Fitting parameter  $P$  and model validation.

**Figure S5.** Fitting parameter  $P$  and model validation: Baffin Bay 2006.

**Figure S6.** Fitting parameter  $P$  and model validation: Baffin Bay 2008.

**Figure S7.** Fitting parameter  $P$  and model validation: Beaufort Sea 2010.

**Figure S8.** Fitting parameter  $P$  and model validation: Beaufort Sea 2011.

**Figure S9.** Fitting parameter  $P$  and model validation: Beaufort Sea 2014.

**Figure S10.** Fitting parameter  $P$  and model validation: Kitikmeot 2006.

**Figure S11.** Fitting parameter  $P$  and model validation: Kitikmeot 2011.

**Figure S12.** Fitting parameter  $P$  and model validation: Kitikmeot 2015.

**Figure S13.** Fitting parameter  $P$  and model validation: Laptev Sea 2003.

**Figure S14.** Fitting parameter  $P$  and model validation: Laptev Sea 2007.

**Table S1.** Model efficiency in predicting length-at-age in 2005 for different larval sizes at hatch.

**Text S2.** Sensitivity of growth to individual model parameters.

**Table S2.** Variation in model parameters and contribution to prediction error ( $R^2$ ) shown by two larval sizes.

**Figure S15.** Sensitivity analysis by individual parameter perturbation (IPP).

**Text S3.** Sensitivity of growth to the combined effect of model parameters.

**Figure S16.** Pearson correlation between modeled larval length (small larvae of 6 mm) and model parameters.

**Figure S17.** Pearson correlation between modeled larval length (large larvae of 12 mm) and model parameters.

**Figure S18.** Parameters ranking from a global sensitivity analysis.

### Funding

This research was undertaken thanks in part to funding from the Canada First Research Excellence Fund, through the Ocean Frontier Institute.

### Competing interests

Authors declare no competing interests.

### Author contributions

- Contributed to conception and design: CLD, RJ, JH.
- Contributed to acquisition of data: CLD, CB.
- Contributed to analysis and interpretation of data: CLD, CB.
- Drafted and/or revised this article: CLD, RJ, CB, HH, JH.
- Approved the submitted version for publication: CLD, RJ, CB, HH, JH.

### References

- Aronovich, TM, Doroshev, SI, Spectorova, LV, Makhotin, VM.** 1975. Egg incubation and larval rearing of navaga (*Eleginus navaga* Pall.), polar cod (*Boreogadus saida* lepechin) and arctic flounder (*Liopsetta glacialis* Pall.) in the laboratory. *Aquaculture* **6**(3): 233–242. DOI: [http://dx.doi.org/10.1016/0044-8486\(75\)90043-5](http://dx.doi.org/10.1016/0044-8486(75)90043-5).
- Bartell, SM, Breck, JE, Gardner, RH, Brenkert, AL.** 1986. Individual parameter perturbation and error analysis of fish bioenergetics models. *Canadian Journal of Fisheries and Aquatic Sciences* **43**(1): 160–168. DOI: <http://dx.doi.org/10.1139/f86-018>.
- Bender, ML, Giebichenstein, J, Teisrud, RN, Laurent, J, Frantzen, M, Meador, JP, Sørensen, L, Hansen, BH, Reinardy, HC, Laurel, B, Nahrgang, J.** 2021. Combined effects of crude oil exposure and warming on eggs and larvae of an arctic forage fish. *Scientific Reports* **11**(1): 8410. DOI: <http://dx.doi.org/10.1038/s41598-021-87932-2>.
- Björnsson, B.** 1999. Is the growth rate of Icelandic cod food-limited? A comparison between pen-reared cod and wild cod living under similar thermal conditions. *Rit Fiskideildar* **16**: 271–279.
- Bouchard, C, Fortier, L.** 2008. Effects of polynyas on the hatching season, early growth and survival of polar cod *Boreogadus saida* in the Laptev Sea. *Marine Ecology Progress Series* **355**: 247–256. DOI: <http://dx.doi.org/10.3354/meps07335>.
- Bouchard, C, Fortier, L.** 2011. Circum-arctic comparison of the hatching season of polar cod *Boreogadus saida*: A test of the freshwater winter refuge hypothesis. *Progress in Oceanography* **90**(1–4):

- 105–116. DOI: <http://dx.doi.org/10.1016/j.pocan.2011.02.008>.
- Bouchard, C, Fortier, L.** 2020. The importance of *Calanus glacialis* for the feeding success of young polar cod: A circumpolar synthesis. *Polar Biology* **43**(8): 1095–1107. DOI: <http://dx.doi.org/10.1007/s00300-020-02643-0>.
- Bouchard, C, Geoffroy, M, LeBlanc, M, Majewski, A, Gauthier, S, Walkusz, W, Reist, JD, Fortier, L.** 2017. Climate warming enhances polar cod recruitment, at least transiently. *Progress in Oceanography* **156**: 121–129. DOI: <http://dx.doi.org/10.1016/j.pocan.2017.06.008>.
- Brett, JR.** 1979. Environmental factors and growth, in Hoar, W, Randall, D, Brett, J, eds., *Fish physiology*. New York, NY: Academic Press. DOI: [http://dx.doi.org/10.1016/S1546-5098\(08\)60033-3](http://dx.doi.org/10.1016/S1546-5098(08)60033-3).
- Brett, JR, Groves, TDD.** 1979. Physiological energetics, in Hoar, W, Randall, D, Brett, J, eds., *Fish physiology*. New York, NY: Academic Press. DOI: [http://dx.doi.org/10.1016/S1546-5098\(08\)60029-1](http://dx.doi.org/10.1016/S1546-5098(08)60029-1).
- Clarke, A.** 2006. Temperature and the metabolic theory of ecology. *Functional Ecology* **20**(2): 405–412. DOI: <http://dx.doi.org/10.1111/j.1365-2435.2006.01109.x>.
- Clarke, A, North, AW.** 1991. Is the growth of polar fish limited by temperature? in di Prisco, G, Maresca, BBT, eds., *Biology of Antarctic fish*. Berlin, Germany: Springer. DOI: [http://dx.doi.org/10.1007/978-3-642-76217-8\\_4](http://dx.doi.org/10.1007/978-3-642-76217-8_4).
- Comita, GW, Marshall, SM, Orr, AP.** 1966. On the biology of *Calanus finmarchicus*: XIII. Seasonal change in weight, calorific value and organic matter. *Journal of the Marine Biological Association of the United Kingdom* **46**(1): 1–17. DOI: <http://dx.doi.org/10.1017/S0025315400017513>.
- Copeman, L, Spencer, M, Heintz, R, Vollenweider, J, Sremba, A, Helser, T, Logerwell, L, Sousa, L, Danielson, S, Pinchuk, AI, Laurel, B.** 2020. Ontogenetic patterns in lipid and fatty acid biomarkers of juvenile polar cod (*Boreogadus saida*) and saffron cod (*Eleginus gracilis*) from across the Alaska Arctic. *Polar Biology* **43**(8): 1121–1140. DOI: <http://dx.doi.org/10.1007/s00300-020-02648-9>.
- Drost, HE, Carmack, EC, Farrell, AP.** 2014. Upper thermal limits of cardiac function for Arctic cod *Boreogadus saida*, a key food web fish species in the Arctic Ocean. *Journal of Fish Biology* **84**(6): 1781–1792. DOI: <http://dx.doi.org/10.1111/jfb.12397>.
- Drost, HE, Fisher, J, Randall, F, Kent, D, Carmack, EC, Farrell, AP.** 2016. Upper thermal limits of the hearts of Arctic cod *Boreogadus saida*: Adults compared with larvae. *Journal of Fish Biology* **88**(2): 718–726. DOI: <http://dx.doi.org/10.1111/jfb.12807>.
- Eriksen, E, Huserbråten, M, Gjøsæter, H, Vikebø, F, Albretsen, J.** 2020. Polar cod egg and larval drift patterns in the Svalbard archipelago. *Polar Biology* **43**: 1–14. DOI: <http://dx.doi.org/10.1007/s00300-019-02549-6>.
- Fiksen, Ø, Jørgensen, C.** 2011. Model of optimal behaviour in fish larvae predicts that food availability determines survival, but not growth. *Marine Ecology Progress Series* **432**: 207–219. DOI: <http://dx.doi.org/10.3354/meps09148>.
- Fiksen, Ø, MacKenzie, B.** 2002. Process-based models of feeding and prey selection in larval fish. *Marine Ecology Progress Series* **243**: 151–164. DOI: <http://dx.doi.org/10.3354/meps243151>.
- Finn, R, Rønnestad, I, van der Meeren, T, Fyhn, H.** 2002. Fuel and metabolic scaling during the early life stages of Atlantic cod *Gadus morhua*. *Marine Ecology Progress Series* **243**: 217–234. DOI: <http://dx.doi.org/10.3354/meps243217>.
- Folkvord, A.** 2005. Comparison of size-at-age of larval Atlantic cod (*Gadus morhua*) from different populations based on size- and temperature-dependent growth models. *Canadian Journal of Fisheries and Aquatic Sciences* **62**(5): 1037–1052. DOI: <http://dx.doi.org/10.1139/f05-008>.
- Geoffroy, M, Majewski, A, LeBlanc, M, Gauthier, S, Walkusz, W, Reist, JD, Fortier, L.** 2016. Vertical segregation of age-0 and age-1+ polar cod (*Boreogadus saida*) over the annual cycle in the Canadian Beaufort Sea. *Polar Biology* **39**(6): 1023–1037. DOI: <http://dx.doi.org/10.1007/s00300-015-1811-z>.
- Graham, M, Hop, H.** 1995. Aspects of reproduction and larval biology of Arctic cod (*Boreogadus saida*). *Arctic* **48**(2): 130–135.
- Holeton, GF.** 1974. Metabolic cold adaptation of polar fish: Fact or artefact? *Physiological Zoology* **47**(3): 137–152. DOI: <http://dx.doi.org/10.1086/physzool.47.3.30157851>.
- Hop, H, Graham, M.** 1995. Respiration of juvenile Arctic cod (*Boreogadus saida*): Effects of acclimation, temperature, and food intake. *Polar Biology* **15**: 359–367.
- Hop, H, Tonn, WM.** 1998. Gastric evacuation rates and daily rations of Arctic cod (*Boreogadus saida*) at low temperatures. *Polar Biology* **19**: 293–301.
- Hop, H, Tonn, WM, Welch, HE.** 1997. Bioenergetics of Arctic cod (*Boreogadus saida*) at low temperatures. *Canadian Journal of Fisheries and Aquatic Sciences* **54**: 1772–1784.
- Ikeda, T.** 2016. Routine metabolic rates of pelagic marine fishes and cephalopods as a function of body mass, habitat temperature and habitat depth. *Journal of Experimental Marine Biology and Ecology* **480**: 74–86. DOI: <http://dx.doi.org/10.1016/j.jembe.2016.03.012>.
- Jørgensen, C, Enberg, K, Mangel, M.** 2016. Modelling and interpreting fish bioenergetics: A role for behaviour, life-history traits and survival trade-offs. *Journal of Fish Biology* **88**(1): 389–402. DOI: <http://dx.doi.org/10.1111/jfb.12834>.
- Kent, D, Drost, HE, Fisher, J, Oyama, T, Farrell, AP.** 2016. Laboratory rearing of wild Arctic cod *Boreogadus saida* from egg to adulthood. *Journal of Fish Biology* **88**(3): 1241–1248. DOI: <http://dx.doi.org/10.1111/jfb.12893>.



- Kitchell, JF, Breck, JE.** 1980. Bioenergetics model and foraging hypothesis for Sea lamprey (*Petromyzon marinus*). *Canadian Journal of Fisheries and Aquatic Sciences* **37**(11): 2159–2168. DOI: <http://dx.doi.org/10.1139/f80-258>.
- Koenker, BL, Laurel, BJ, Copeman, LA, Ciannelli, L.** 2018. Effects of temperature and food availability on the survival and growth of larval Arctic cod (*Boreogadus saida*) and walleye pollock (*Gadus chalcogrammus*). *ICES Journal of Marine Science* **75**(7): 2386–2402. DOI: <http://dx.doi.org/10.1093/icesjms/fsy062>.
- Kohlbach, D, Graeve, MA, Lange, B, David, C, Peeken, I, Flores, H** 2016. The importance of ice algae-produced carbon in the central Arctic Ocean ecosystem: Food web relationships revealed by lipid and stable isotope analyses. *Limnology and Oceanography* **61**(6): 2027–2044. DOI: <http://dx.doi.org/10.1002/lno.10351>.
- Kohlbach, D, Schaafsma, FL, Graeve, M, Lebreton, B, Lange, BA, David, C, Vortkamp, M, Flores, H.** 2017. Strong linkage of polar cod (*Boreogadus saida*) to sea ice algae-produced carbon: Evidence from stomach content, fatty acid and stable isotope analyses. *Progress in Oceanography* **152**: 62–74. DOI: <http://dx.doi.org/10.1016/j.pocean.2017.02.003>.
- Kristiansen, T, Fiksen, Ø, Folkvord, A.** 2007. Modelling feeding, growth, and habitat selection in larval Atlantic cod (*Gadus morhua*): Observations and model predictions in a macrocosm environment. *Canadian Journal of Fisheries and Aquatic Sciences* **64**(1): 136–151. DOI: <http://dx.doi.org/10.1139/f06-176>.
- Kristiansen, T, Jorgensen, C, Lough, RG, Vikebo, F, Fiksen, O.** 2009. Modeling rule-based behavior: Habitat selection and the growth-survival trade-off in larval cod. *Behavioral Ecology* **20**(3): 490–500. DOI: <http://dx.doi.org/10.1093/beheco/arp023>.
- Kunz, KL, Frickenhaus, S, Hardenberg, S, Johansen, T, Leo, E, Pörtner, HO, Schmidt, M, Windisch, HS, Knust, R, Mark, FC.** 2016. New encounters in Arctic waters: A comparison of metabolism and performance of polar cod (*Boreogadus saida*) and Atlantic cod (*Gadus morhua*) under ocean acidification and warming. *Polar Biology* **39**(6): 1137–1153. DOI: <http://dx.doi.org/10.1007/s00300-016-1932-z>.
- Laurel, BJ, Copeman, LA, Spencer, M, Iseri, P.** 2017. Temperature-dependent growth as a function of size and age in juvenile Arctic cod (*Boreogadus saida*). *ICES Journal of Marine Science* **74**(6): 1614–1621. DOI: <http://dx.doi.org/10.1093/icesjms/fsx028>.
- Laurel, BJ, Copeman, LA, Spencer, M, Iseri, P.** 2018. Comparative effects of temperature on rates of development and survival of eggs and yolk-sac larvae of Arctic cod (*Boreogadus saida*) and walleye pollock (*Gadus chalcogrammus*). *ICES Journal of Marine Science* **75**(7): 2403–2412. DOI: <http://dx.doi.org/10.1093/icesjms/fsy042>.
- Laurel, BJ, Spencer, M, Iseri, P, Copeman, LA.** 2016. Temperature-dependent growth and behavior of juvenile Arctic cod (*Boreogadus saida*) and co-occurring North Pacific gadids. *Polar Biology* **39**(6): 1127–1135. DOI: <http://dx.doi.org/10.1007/s00300-015-1761-5>.
- LeBlanc, M, Geoffroy, M, Bouchard, C, Gauthier, S, Majewski, A, Reist, JD, Fortier, L.** 2020. Pelagic production and the recruitment of juvenile polar cod *Boreogadus saida* in Canadian Arctic seas. *Polar Biology* **43**: 1043–1054. DOI: <http://dx.doi.org/10.1007/s00300-019-02565-6>.
- Lough, RG, Buckley, LJ, Werner, FE, Quinlan, JA, Pehrson Edwards, K.** 2005. A general biophysical model of larval cod (*Gadus morhua*) growth applied to populations on Georges Bank. *Fisheries Oceanography* **14**(4): 241–262. DOI: <http://dx.doi.org/10.1111/j.1365-2419.2005.00330.x>.
- Mauchline, J.** 1998. *The biology of calanoid copepods*. London, UK: Academic Press. (Advances in marine biology; vol. 33). ISBN 0-12-026133-2.
- Mayer, DG, Butler, DG.** 1993. Statistical validation. *Ecological Modelling* **68**(1–2): 21–32. DOI: [http://dx.doi.org/10.1016/0304-3800\(93\)90105-2](http://dx.doi.org/10.1016/0304-3800(93)90105-2).
- Mecklenburg, C, Steinke, D.** 2015. Ichthyofaunal baselines in the Pacific Arctic region and RUSALCA study area. *Oceanography* **28**(3): 158–189. DOI: <http://dx.doi.org/10.5670/oceanog.2015.64>.
- Menemenlis, D, Campin, J, Heimbach, P, Hill, C, Lee, T, Nguyen, A, Schodlok, M, Zhang, H.** 2008. ECCO2: High resolution global ocean and sea ice data synthesis. *Mercator Ocean Quarterly Newsletter* **31**: 13–21. Available at [http://apdr.csoest.hawaii.edu/erddap/griddap/hawaii\\_soest\\_daa1\\_6006\\_642b.html](http://apdr.csoest.hawaii.edu/erddap/griddap/hawaii_soest_daa1_6006_642b.html). Accessed 7 May 2021.
- Michaud, J, Fortier, L, Rowe, P, Ramseier, R.** 1996. Feeding success and survivorship of Arctic cod larvae, *Boreogadus saida*, in the Northeast Water polynya (Greenland Sea). *Fisheries Oceanography* **5**(2): 120–135. DOI: <http://dx.doi.org/10.1111/j.1365-2419.1996.tb00111.x>.
- Monk, J, Puvanendran, V, Brown, JA.** 2006. Do different light regimes affect the foraging behaviour, growth and survival of larval cod (*Gadus morhua* L.)? *Aquaculture* **257**(1–4): 287–293. DOI: <http://dx.doi.org/10.1016/j.aquaculture.2006.02.071>.
- Mueter, F, Bouchard, C, Hop, H, Laurel, B, Norcross, B.** 2020. Arctic gadids in a rapidly changing environment. *Polar Biology* **43**: 945–949. DOI: <http://dx.doi.org/10.1007/s00300-020-02696-1>.
- Munk, P.** 1997. Prey size spectra and prey availability of larval and small juvenile cod. *Journal of Fish Biology* **51**(sa): 340–351. DOI: <http://dx.doi.org/10.1111/j.1095-8649.1997.tb06107.x>.
- Otterlei, E, Nyhammer, G, Folkvord, A, Stefansson, SO.** 1999. Temperature- and size-dependent growth of larval and early juvenile Atlantic cod (*Gadus morhua*): A comparative study of Norwegian coastal cod and northeast Arctic cod. *Canadian Journal of Fisheries and Aquatic Sciences* **56**(11): 2099–2111. DOI: <http://dx.doi.org/10.1139/f99-168>.

- Payne, MF, Rippingale, RJ.** 2000. Rearing West Australian seahorse, *Hippocampus subelongatus*, Juveniles on copepod nauplii and enriched *Artemia*. *Aquaculture* **188**(3–4): 353–361. DOI: [http://dx.doi.org/10.1016/S0044-8486\(00\)00349-5](http://dx.doi.org/10.1016/S0044-8486(00)00349-5).
- Peck, MA, Daewel, U.** 2007. Physiologically based limits to food consumption, and individual-based modeling of foraging and growth of larval fishes. *Marine Ecology Progress Series* **347**: 171–183. DOI: <http://dx.doi.org/10.3354/meps06976>.
- Peck, MA, Hufnagl, M.** 2012. Can IBMs tell us why most larvae die in the sea? Model sensitivities and scenarios reveal research needs. *Journal of Marine Systems* **93**: 77–93. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2011.08.005>.
- Pepin, P, Robert, D, Bouchard, C, Dower, JF, Falardeau, M, Fortier, L, Jenkins, GP, Leclerc, V, Levesque, K, Llopiz, JK, Meekan, MG.** 2014. Once upon a larva: Revisiting the relationship between feeding success and growth in fish larvae. *ICES Journal of Marine Science* **72**(2): 359–373. DOI: <http://dx.doi.org/10.1093/icesjms/fsu201>.
- Ponomarenko, VP.** 2000. Eggs, larvae, and juveniles of polar cod *Boreogadus saida* in the Barents, Kara, and White Seas. *Journal of Ichthyology* **40**: 165–173.
- Pörtner, H-O, Storch, D, Heilmayer, O.** 2005. Constraints and trade-offs in climate-dependent adaptation: Energy budgets and growth in a latitudinal cline. *Scientia Marina* **69**(Suppl. 2): 271–285. DOI: <http://dx.doi.org/10.3989/scimar.2005.69s2271>.
- Puvanendran, V, Brown, JA.** 1999. Foraging, growth and survival of Atlantic cod larvae reared in different prey concentrations. *Aquaculture* **175**(1–2): 77–92. DOI: [http://dx.doi.org/10.1016/S0044-8486\(99\)00023-X](http://dx.doi.org/10.1016/S0044-8486(99)00023-X).
- Railsback, SF, Rose, KA.** 1999. Bioenergetics modeling of stream trout growth: Temperature and food consumption effects. *Transactions of the American Fisheries Society* **128**(2): 241–256. DOI: [http://dx.doi.org/10.1577/1548-8659\(1999\)128<0241:bmostg>2.0.co;2](http://dx.doi.org/10.1577/1548-8659(1999)128<0241:bmostg>2.0.co;2).
- Ricker, WE.** 1979. Growth rates and models, in Hoar, W, Randall, D, Brett, J, eds., *Fish physiology*. New York, NY: Academic Press. DOI: [http://dx.doi.org/10.1016/S1546-5098\(08\)60034-5](http://dx.doi.org/10.1016/S1546-5098(08)60034-5).
- Roy, D, Haffner, GD, Brandt, SB.** 2004. Estimating fish production potentials using a temporally explicit model. *Ecological Modelling* **173**(2–3): 241–257. DOI: <http://dx.doi.org/10.1016/j.ecolmodel.2003.06.005>.
- Sakurai, Y, Ishii, K, Nakatani, T, Yamaguchi, H, Anma, G, Jin, M.** 1998. Reproductive characteristics and effects of temperature and salinity on the development and survival of eggs and larvae of Arctic cod (*Boreogadus saida*). *Memoirs of the Faculty of Fisheries Hokkaido University* **48**(1): 77–89. Available at <https://eprints.lib.hokudai.ac.jp/dspace/handle/2115/21924>.
- Shields, RJ, Bell, JG, Luizi, FS, Gara, B, Bromage, NR, Sargent, JR.** 1999. Natural copepods are superior to enriched *Artemia* nauplii as feed for halibut larvae (*Hippoglossus hippoglossus*) in terms of survival, pigmentation and retinal morphology: Relation to dietary essential fatty acids. *Journal of Nutrition* **129**(6): 1186–1194. DOI: <http://dx.doi.org/10.1093/jn/129.6.1186>.
- Sogard, SM.** 1997. Size-selective mortality in the juvenile stage of teleost fishes: A review. *Bulletin of Marine Science* **60**: 1129–1157.
- Sorgeloos, P, Dhert, P, Candreva, P.** 2001. Use of the brine shrimp, *Artemia* spp., in marine fish larviculture. *Aquaculture* **200**(1–2): 147–159. DOI: [http://dx.doi.org/10.1016/S0044-8486\(01\)00698-6](http://dx.doi.org/10.1016/S0044-8486(01)00698-6).
- Støttrup, JG.** 2000. The elusive copepods: Their production and suitability in marine aquaculture. *Aquaculture Research* **31**(8–9): 703–711. DOI: <http://dx.doi.org/10.1046/j.1365-2109.2000.318488.x>.
- Thanassekos, S, Fortier, L.** 2012. An individual based model of Arctic cod (*Boreogadus saida*) early life in Arctic polynyas : I. Simulated growth in relation to hatch date in the Northeast Water (Greenland Sea) and the North Water (Baffin Bay). *Journal of Marine Systems* **93**: 25–38. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2011.08.003>.
- Thanassekos, S, Robert, D, Fortier, L.** 2012. An individual based model of Arctic cod (*Boreogadus saida*) early life in Arctic polynyas: II. Length-dependent and growth-dependent mortality. *Journal of Marine Systems* **93**: 39–46. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2011.08.001>.
- van Griensven, A, Meixner, T, Grunwald, S, Bishop, T, Diluzio, M, Srinivasan, R.** 2006. A global sensitivity analysis tool for the parameters of multi-variable catchment models. *Journal of Hydrology* **324**(1–4): 10–23. DOI: <http://dx.doi.org/10.1016/j.jhydrol.2005.09.008>.
- Vanhaecke, P, Lavens, P, Sorgeloos, P.** 1983. International study on *Artemia*. XVII. Energy consumption in cysts and early larval stages of various geographical strains of *Artemia*. *Annales de la Société royale zoologique de Belgique* **113**(2): 155–164.
- Vestfals, CD, Mueter, FJ, Hedstrom, KS, Laurel, BJ, Petrik, CM, Duffy-Anderson, JT, Danielson, SL.** 2021. Modeling the dispersal of polar cod (*Boreogadus saida*) and saffron cod (*Eleginus gracilis*) early life stages in the Pacific Arctic using a biophysical transport model. *Progress in Oceanography* **196**: 102571. DOI: <http://dx.doi.org/10.1016/j.pocean.2021.102571>.
- Walkusz, W, Paulic, JE, Williams, WJ, Kwasniewski, S, Papst, MH.** 2011. Distribution and diet of larval and juvenile Arctic cod (*Boreogadus saida*) in the shallow Canadian Beaufort Sea. *Journal of Marine Systems* **84**(3–4): 78–84. DOI: <http://dx.doi.org/10.1016/j.jmarsys.2010.09.001>.

**How to cite this article:** David, CL, Ji, R, Bouchard, C, Hop, H, Hutchings, JA. 2022. The interactive effects of temperature and food consumption on growth of larval Arctic cod (*Boreogadus saida*): A bioenergetic model. *Elementa: Science of the Anthropocene* 10(1). DOI: <https://doi.org/10.1525/elementa.2021.00045>

**Domain Editor-in-Chief:** Jody W. Deming, University of Washington, Seattle, WA, USA

**Guest Editor:** Dominique Robert, Institut des Sciences de la Mer, Université du Québec à Rimouski, Canada

**Knowledge Domain:** Ocean Science

**Part of an Elementa Special Feature:** Four Decades of Arctic Climate Change: A Tribute to Louis Fortier

**Published:** February 10, 2022    **Accepted:** January 17, 2022    **Submitted:** June 11, 2021

**Copyright:** © 2022 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See <http://creativecommons.org/licenses/by/4.0/>.



*Elem Sci Anth* is a peer-reviewed open access journal published by University of California Press.

OPEN ACCESS