



# Does an obesogenic family environment moderate the association between sports participation and body composition in children? The ENERGY project

Mette Stavnsbo<sup>1</sup>  | Andreas Stenling<sup>1,2</sup> | Sveinung Berntsen<sup>1</sup> |  
 Mai J. M. Chinapaw<sup>3,4</sup>  | Bjørge H. Hansen<sup>1</sup> | Yannis Manios<sup>5,6</sup> |  
 Dénes Molnár<sup>7</sup> | Monica K. Torstveit<sup>1</sup> | Maïté Verloigne<sup>8</sup> |  
 Germán Vicente-Rodríguez<sup>9</sup> | Thomas Westergren<sup>10,11</sup> | Elling Bere<sup>1,11</sup>

<sup>1</sup>Department of Sports Science and Physical Education, University of Agder, Kristiansand, Norway

<sup>2</sup>Department of Psychology, Umeå University, Umeå, Sweden

<sup>3</sup>Amsterdam UMC, location Vrije Universiteit Amsterdam, Public and Occupational Health, Amsterdam, The Netherlands

<sup>4</sup>Amsterdam Public Health, Health Behaviour and Chronic Diseases and Methodology, Amsterdam, The Netherlands

<sup>5</sup>Department of Nutrition & Dietetics, School of Health Sciences & Education, Harokopio University, Athens, Greece

<sup>6</sup>Institute of Agri-food and Life Sciences, Hellenic Mediterranean University Research Centre, Heraklion, Greece

<sup>7</sup>Department of Pediatrics, Medical School, University of Pécs, Pécs, Hungary

<sup>8</sup>Department of Public Health and Primary Care, Ghent University, Ghent, Belgium

<sup>9</sup>GENUD (Growth, Exercise, NUtrition and Development) Research Group, FIMS Collaborating Center of Sports Medicine, Instituto Agroalimentario de Aragón—IA2—(CITA-Universidad de Zaragoza), Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición (CIBEROBn), Faculty of Health and Sport Science, Department of Psychiatry and Nursing, University of Zaragoza, Zaragoza, Spain

<sup>10</sup>Department of Health and Nursing Science, University of Agder, Kristiansand, Norway

<sup>11</sup>Department of Health and Inequalities & Centre for Evaluation of Public Health Measures, Norwegian Institute of Public Health, Oslo, Norway

## Correspondence

Mette Stavnsbo, Department of Sport Science and Physical Education, University of Agder, PO Box 422, N-4604 Kristiansand, Norway.  
 Email: [mette.stavnsbo@uia.no](mailto:mette.stavnsbo@uia.no)

## Funding information

FP7 Ideas: European Research Council, Grant/Award Number: 223254; Netherlands Organization for Health Research and Development, Grant/Award Number: 50-50150-98-002

## Summary

**Background:** Previous research on the association between sports participation and body composition has shown mixed findings. The family home is considered one of the most influential environments on childhood obesity. Thus, the association between sports participation and body composition in children may be influenced by an obesogenic home environment.

**Objectives:** To investigate if an obesogenic family environment moderates the association between sports participation and body composition in children.

**Methods:** A total of 3999 children (54% girls; 11.6 ± 0.7 years) and their parent(s) were included from the ENERGY project. A composite obesogenic family environment risk score was created from 10 questionnaire items. Height, weight (to calculate body mass index), and waist circumference were obtained by trained researchers and used as indicators of body composition.

**Abbreviations:** BMI, body mass index; CI, confidence interval; ICC, intraclass correlation coefficients; MVPA, moderate-to-vigorous PA; PA, physical activity; SD, standard deviation; SES, socio-economic status; WC, waist circumference.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Pediatric Obesity* published by John Wiley & Sons Ltd on behalf of World Obesity Federation.

**Results:** The composite risk score significantly moderated the association between sports participation and both waist circumference and body mass index. In children from families with moderate and high obesogenic risk, organized sports participation was significantly associated with smaller waist circumference (moderate risk:  $-0.29$ , 95% CI  $-0.45$  to  $-0.14$ ; high risk:  $-0.46$ , 95% CI  $-0.66$  to  $-0.25$ ) and lower body mass index (moderate risk:  $-0.10$ , 95% CI  $-0.16$  to  $-0.04$ ; high risk:  $-0.14$ , 95% CI  $-0.22$  to  $-0.06$ ), but not in children with a low obesogenic family risk score.

**Conclusions:** Enrolling children in sports activities from an early age can be important for healthy weight maintenance, especially among children from obesogenic family environments.

#### KEYWORDS

obesity, obesogenic environment, overweight, paediatrics, physical activity

## 1 | INTRODUCTION

Paediatric obesity is a global public health concern associated with a myriad of adverse health consequences.<sup>1</sup> Physical activity (PA) is important for the maintenance of a healthy weight, in addition to other health benefits.<sup>2</sup> Leisure time sport is one component of PA that can contribute to public health. For instance, children and adolescents participating in sporting clubs are more likely to meet overall PA recommendations of 60 min of moderate-to-vigorous intensity PA (MVPA) daily.<sup>3</sup> Thus, sporting clubs are an important venue for PA and health promotion. Some studies report higher sports participation to be associated with healthier indicators of body composition, such as body mass index (BMI) and waist circumference (WC) among children (i.e., from the ENERGY project).<sup>4</sup> In general, however, the evidence is mixed. In a systematic review by Lee et al.,<sup>5</sup> six out of 17 studies found no statistically significant associations between sports participation and child obesity or only partial or weak associations among subgroups of children. Despite these inconsistent findings, PA is hypothesized to protect against the development of obesity resulting in higher energy expenditure and resting metabolic rate.<sup>6</sup> Obesity is, nevertheless, both a complex and multifactorial condition.

Socio-ecological models can be helpful to understand the complexity of risk factors contributing to childhood obesity.<sup>7,8</sup> Especially among younger children, the home (or family) environment is considered the most important influencing factor.<sup>7</sup> Although there is no consensus regarding the exact risk factors an obesogenic family environment encompasses, researchers have highlighted the importance of factors related to PA, food intake, and media use, which can be expressed in both the physical- (i.e., accessibility to foods and media) and social- (i.e., parental role modelling and rules) family environment.<sup>8</sup>

In the European Energy balance Research to prevent excessive weight Gain among Youth (ENERGY) project,<sup>9</sup> data were collected on several physical and social factors in a family environment that have been associated with increased risk of child obesity. Findings from the ENERGY project show that families dining together and parenting rules regarding media use (e.g., not allowing TV while eating and TV in the child's bedroom) are associated with healthier body composition among children.<sup>10-13</sup> Furthermore, parental socioeconomic status (SES) and

BMI,<sup>14,15</sup> in addition to parental behaviours such as daily intake of sugar sweetened drinks, PA levels, and sedentary time,<sup>4,16-18</sup> have been associated with their children's weight-related health behaviours and weight status. Although each of these factors have shown to be associated with weight-related outcomes among children, a composite measure should capture overall obesogenic risk in the family home environment more efficiently than single risk factors.<sup>19</sup>

A better understanding of the mixed findings regarding the associations between child sports participation and body composition is warranted. Some children may benefit more from increased PA and sports participation than others, for instance, children with multiple unhealthy lifestyle behaviours.<sup>20</sup> This reasoning is supported by intervention studies showing that increased sports participation particularly seem to benefit overweight children living in environments with higher obesogenic risk (e.g., low-income families and obese parents).<sup>21,22</sup> Based on previous research it is reasonable to assume that different risk (i.e., obesogenic family environment risk) and protective factors (i.e., sports participation) may interact in their influence on child body composition.<sup>23</sup> More specifically, the protective effect of sports participation against obesity might be more pronounced among children with an increased risk of developing excessive weight due to an obesogenic family environment. Therefore, the present study aims to examine whether a composite obesogenic family environmental risk score, based on factors related to both the physical and social environment in the family, moderates the association between sports participation and body composition in 10-to-12-year-old children across Europe. We hypothesize that the association between sports participation and body composition is stronger in children from high obesogenic risk environments compared to children from low obesogenic risk environments.

## 2 | METHODS

### 2.1 | Procedure and sample

The current study used data from the ENERGY (European Energy balance Research to prevent excessive weight Gain among Youth)

project, more specifically from the cross-sectional study conducted in seven European countries (Belgium, Greece, Hungary, The Netherlands, Norway, Slovenia, and Spain).<sup>9</sup> Each country was represented by a local research institute or university department, being responsible for the data collection in the respective country. Ethical approval was obtained in all seven participating countries from Medical Ethical review committees and all participants and legal guardians provided informed consent prior to commencement of investigation. The standardized sampling procedures, data collection, data handling, and ethical approval in the separate countries are described in detail elsewhere.<sup>9</sup> Briefly, the cross-sectional survey was carried out in primary schools among 10–12-year-old children. The study aimed to include a minimum sample of 1000 school children from each participating country and one parent/caretaker for each child. The recruitment and data collection took place from March–July 2010. Nationally representative samples were obtained from Greece, Hungary, The Netherlands, and Slovenia. In Belgium, schools were selected from Flanders (i.e., the northern Dutch-speaking part of Belgium), Norway selected schools from the southern regions of the country, and Spain selected schools in the region of Aragón. Between 15 (Slovenia) and 37 (Greece) schools participated, with a response rate at the school level between 5% (The Netherlands) and 100% (Slovenia). Response rates were in general high at the child level (<85%); however, lower in Hungary (33%) and Norway (45%), primarily due to a lack of completed parental consent forms. At the parental level, general response rates were also relatively high (80%).

In total, 7234 children and 6002 parents completed the questionnaires. Children completed the child questionnaire during class-time with guidance from a research assistant or project worker, and each child brought home a parental questionnaire to be completed by one of the parents.<sup>9</sup>

## 2.2 | Measures

### 2.2.1 | Child sports participation

Children were asked how many hours per week they participated in one or two sports ( $ICC_{\text{test-retest}} = 0.74$  for sport 1; 1.00 for sport 2).<sup>24</sup> Total hours of sport participation per week was calculated for each child.<sup>4</sup>

### 2.2.2 | Child body composition

Body height, weight, and WC were measured by trained research assistants according to standardized protocols ( $ICC_{\text{test-retest}} = 1.00$  for weight and height; 0.99 for WC).<sup>9,24</sup> The children were measured in light clothing without shoes. Body height was measured with a SECA Leicester Portable stadiometer (accuracy of 0.1 cm) and weight measured with a calibrated electronic scale SECA 861 (accuracy of 0.1 kg). Waist circumference was measured with a SECA 201 measuring band (accuracy of 0.1 cm). Two readings of each measurement

(height, weight, and WC) were obtained. A third measure was taken if the two readings differed more than 1%.<sup>9</sup> Body mass index ( $\text{kg}/\text{m}^2$ ) was calculated as weight (kg) divided by height squared ( $\text{m}^2$ ), and weight categories based on the International Obesity Task Force criteria (IOTF).<sup>25</sup>

### 2.2.3 | Ethnicity

Ethnicity was assessed in the child questionnaire and operationalized as a distinction between children of immigrant origin (non-native) and of native origin, based on the primary language spoken at home. The answering categories were tailored to the different countries. This included the official language or languages of the specific country or region, the native languages of the largest ethnic minorities, and a category 'other'. A dichotomous variable was created that distinguished children for whom the official language of the country of administration was the main language spoken at home (native) coded 0, from those where another language was the main language spoken at home (non-native) coded 1.<sup>26</sup>

### 2.2.4 | Parental marital status

Parental marital status was assessed in the parental questionnaire. The answer 'married' or 'living with my partner but not married' was considered a dual marital status (coded 0), and the answers 'single', 'divorced', or 'separated' as single status (coded 1).<sup>9</sup>

### 2.2.5 | Obesogenic family environment variables

#### *Parental education levels*

Educational level was assessed in the parental questionnaire by asking how many years of education both parents had. In line with previous publications from the ENERGY project,<sup>13,26</sup> educational level was dichotomized as follows: if both parents had <14 years of education each, corresponding to high school across countries as the highest education level, parental educational level was determined as low (coded 1), and if at least one parent had >14 years of education, parental educational level was categorized as high (coded 0).

#### *Parental anthropometrics*

Parental BMI was calculated from self-reported weight and height obtained in the parental questionnaire.<sup>9</sup> The BMI variable was dichotomized into normal weight (including underweight) <25 (coded as 0) and overweight/obese  $\geq 25$  (coded as 1) according to cut-offs by the IOTF criteria.<sup>25</sup>

#### *Parental physical activity*

In the parental questionnaire, active commuting by bicycle to work was assessed by asking how many days a week the parent usually biked. Response categories were between 0 to 5 days/week.<sup>9</sup>

Dichotomization was done according to earlier used cut-offs in the ENERGY study for active commuting to work. Four days or more of bicycling to work was considered a regular routine in weekdays and defined as a high mode (coded as 0), while <4 days was considered as a low mode (coded 1).<sup>27</sup>

Parental leisure time PA was assessed separately for weekdays and weekend days by asking how many days the parents usually participated in physical activities/sports in their leisure time. Response categories were between 0 and 5 days for weekdays, and between 0 and 2 days for weekends.<sup>9</sup> Total days of leisure time PA/sports participation per week was calculated. In the lack of an official recommendation or reference to guide our decision of a cut-off point, we based our decision on the median split similar to how other composite risk scores have been created.<sup>28</sup> The central tendency for the sample was 2 days/week, thus, parental PA/sports participation >2 days per week was defined as high (coded as 0), and weekly PA/sports participation ≤2 days defined as low (coded as 1).

#### Parental screen time

Parents were asked about the average time spent daily watching TV (including video and DVD) and time spent on the computer in their leisure time, separately for weekdays and weekend days.<sup>9</sup> Total hours of parental screen time per week (weekday and weekend days) from both TV and computer was calculated by summing up the average reported hours. No official recommendations of total screen time exist, thus the median split was used to categorize parental screen time into low levels (coded 0) if the weekly average was <5 h of total screen time, while ≥5 h of total screentime was categorized into high levels (coded 1).

#### TV during meals and TV in child bedroom

Parents were asked how often they watched TV during meals (breakfast, lunch and dinner) and reported on a 5-point scale from never to always. According to earlier used cut-offs in the ENERGY project the answer 'never' or 'not often' was considered as a low occurrence and recoded as 0, and 'sometimes, often or always' was recoded 1.<sup>10</sup> The parents were also asked if the child had a TV in the bedroom. The answer 'no' was coded 0 and 'yes' coded 1.<sup>9</sup>

#### Parental intake of sugared drinks

Parents were asked how many times a week they drank sweetened beverages and reported on a 7-point scale from never to more than once a day.<sup>9</sup> In lack of official recommendations, the median split was used to categorize the variable into low intake of sugared drinks (coded 0) representing 'never or less than once a week' and high intake of sugared drinks (coded 1) 'once a week, 2–4 days a week, 5–6 days a week, every day (once a day), and every day (more than once a day)'.

#### Eating breakfast together as a family

Parents were asked how often at least one of the parents were eating breakfast together with the child, answering on a 5-point scale from never to everyday. According to earlier used cut-offs in the ENERGY

project, the answers were dichotomized into 5–7 days representing 'every day' (coded 0) and 2–4 days or less as 'not every day' (coded 1).<sup>13</sup>

## 2.3 | Statistical analysis

Descriptive statistics were performed to present the sample characteristics. Results are presented as mean and standard deviation (SD) or numbers and percentage (%). In addition, bivariate correlations coefficients between the single obesogenic family environment variables, WC, BMI, and sports participation can be found in the supplementary material (Table S1).

The composite obesogenic family environment risk score was defined per child by summing up the dichotomized obesogenic family environment variables, where 0 represented 'low risk' and 1 'high risk'. Thus, a composite risk score was generated with a possible score between 0 (lowest risk) and 9 (highest risk). The composite risk score was normally distributed (mean 4.5 (SD ±1.9)).

The analytical sample included children with complete data on each obesogenic family environment variable, covariates, and dependent variables ( $n = 3999$ ). Compared to those excluded, the analytical sample had slightly lower child sports participation (−0.14 h, 95% CI (−0.27 to −0.01),  $p < 0.050$ ), lower child WC (−0.43 cm, 95% CI (−0.85 to −0.02),  $p < 0.050$ ), more often parents with partners/married ( $X^2(1) \geq 801.946$ ,  $v = 0.37$ ,  $p < 0.001$ ), and parents with higher education levels ( $X^2(1) \geq 5.501$ ,  $v = 0.03$ ,  $p < 0.050$ ), whereas no statistically significant differences were found between groups in child BMI, ethnicity, parental PA levels, or parental weight status.

Linear regression analysis was used to explore the association between the main variables of interest. A significant inverse association between sports participation and both WC and BMI have been reported earlier,<sup>4</sup> thus only the association between the obesogenic composite risk score and body composition was presented in this study.

Path analysis and cluster-robust standard error estimation,<sup>29</sup> to account for a potential clustering effect by school, was used to investigate the moderating effect of the composite obesogenic family environment risk score on the association between sports participation and body composition. First, the interaction term (sports participation\*composite risk score) was calculated, after mean centring the variables. Path models included the interaction term *sports participation\*composite risk score* to investigate the moderation effect of the composite obesogenic family environment risk score on the association between sports participation and both measures of body composition (WC and BMI). The school intraclass correlation coefficients were low (all ICC <0.03), in line with what have been reported earlier from the ENERGY study.<sup>4</sup> Thus, sensitivity analysis was performed to investigate the degree of influence by school clustering, comparing models with and without school as a cluster variable. Very small or no changes in coefficients were observed between the two models, thus no cluster adjustment for school was included in the final model (see model output including clustering by school in the

**TABLE 1** Descriptive statistics of the included study sample by sex ( $N = 3999$ ).

	Girls	Boys
$N$ children (%) <sup>a</sup>	2162 (54)	1837 (46)
Age (mean (SD) yrs)	11.6 (0.7)	11.7 (0.7)
Ethnicity (% native)	1765 (81)	1478 (80)
WC (mean (SD) cm)	<b>65.4 (8.4)</b>	<b>67.7 (9.4)</b>
BMI (kg/m <sup>2</sup> )	19.0 (3.3)	19.2 (3.3)
Normal weight (%) <sup>a</sup>	<b>1734 (80)</b>	<b>1353 (74)</b>
Overweight (%) <sup>a</sup>	<b>355 (16)</b>	<b>403 (21)</b>
Obese (%) <sup>a</sup>	73 (3)	81 (4)
Sports participation (% yes) <sup>a</sup>	1553 (71)	1510 (81)
Sports participation (mean (SD)) hours/week)	<b>3.5 (2.7)</b>	<b>4.3 (2.8)</b>
TV on bedroom (% yes)	692 (32)	663 (36)
Composite risk score (mean (SD))	4.6 (1.9)	4.5 (1.9)
<i>Parental measures</i>		
Marital status (% married)	2066 (95)	1774 (95)
Parental education (% with $\geq 14$ years)	1475 (68)	1284 (69)
Parental BMI ((mean SD) kg/m <sup>2</sup> )	24.6 (4.3)	24.5 (4.2)
Overweight/obese (% yes) <sup>a</sup>	840 (39)	701 (38)
Leisure time activity (% yes)	1401 (64)	1250 (67)
Leisure time PA (mean (SD)) days/week)	2.0 (2.0)	2.0 (2.1)
Screen time (mean (SD) hours/day)	5.1 (2.6)	5.1 (2.6)

Note: Numbers in bold represent significant differences between sex ( $p \leq 0.05$ ).

Abbreviations: BMI, body mass index;  $N$ , number; PA, physical activity; SD, standard deviation; WC, waist circumference; yrs, years.

<sup>a</sup> $N$  (%), number (percentage).

supplementary material Table 1). Further, differences between countries in the magnitude of the association(s) was examined using multi-group analysis. Models with these associations forced to equality between countries was compared to models where they were freely estimated. The unconstrained models had lower Bayesian information criteria (BIC) values than the constrained models, thus indicating better model fit. However, since the difference in BIC values between the constrained and unconstrained models were very small (BIC difference  $< 1\%$ ),<sup>30</sup> path models were run with and without country as a fixed effect to further investigate the influence by country on these associations. The sensitivity analyses confirmed the small differences in BIC values and country was excluded from the final models (see output for models adjusting for country in Table S2). Thus, the final path models, did not adjust for clustering within school and country. If a statistically significant moderation was observed by the composite obesogenic family risk score, the difference in magnitude of the association between sports participation and body composition variable was further explored in tertiles (low, moderate, and high composite risk score). *Second*, regions of significance were calculated using the Johnson-Neyman technique<sup>31</sup> to derive points along the continuum of the moderator where the regression coefficients of the predictor

transitions between statistically significant and non-significant. All analyses were adjusted for age, sex, ethnicity of the child, and parental marital status that earlier have shown an association with sports participation and body composition.<sup>32–34</sup> Results are presented with beta-coefficients and 95% confidence intervals (95% CI).

An alpha-level of  $p \leq 0.05$  was considered statistically significant. IBM SPSS version 23 (IBM SPSS Statistics for Windows, Armonk, NY: IBM Corp.) was used to perform the descriptive statistics, whereas the main analyses were conducted using MPlus version 8 (Muthén and Muthén, 1998–2017).

## 3 | RESULTS

### 3.1 | Descriptive statistics

Out of the total of 7234 children (52% girls;  $11.6 \pm 0.7$  years) that participated in the ENERGY school-based survey, 3999 children (54% girls;  $11.6 \pm 0.7$  years) had valid data on all variables in the main analysis and thus eligible for the present study.

Compared to girls, boys had higher WC and were more often categorized as overweight according to their BMI; however, boys spent more time participating in sports than did girls (Table 1), as also reported earlier in the ENERGY project.<sup>4</sup>

### 3.2 | Association analyses

A statistically significant positive association was found between the composite obesogenic family environment risk score and both WC ( $\beta$  1.11, 95% CI 0.97 to 1.25,  $p < 0.001$ ) and BMI ( $\beta$  0.36, 95% CI 0.31 to 0.41,  $p < 0.001$ ) (Table S3). For every one unit increase in the composite risk score, WC increased by 1.11 cm and BMI by 0.36.

The moderation analyses showed that the association between sports participation and WC ( $-0.10$ , 95% CI  $-0.15$  to  $-0.05$ ),  $p < 0.001$ ) as well as the association between sport participation and BMI ( $-0.04$ , 95% CI  $-0.06$  to  $-0.02$ ),  $p < 0.001$ ) were moderated by the composite risk score. The beta-coefficients for the interaction terms are the difference in 'effect' of sports participation on body composition per one unit increase in the composite score. Thus, the inverse association between sports participation and body composition was stronger in participants with a higher composite obesogenic family environment risk score.

To further examine these associations, we created tertiles of the composite risk score and examined the association between sports participation and body composition within each tertile (Table 2, see also Figures S1 and S2 in the supporting information). Neither the association between sports participation and WC nor the association between sports participation and BMI was statistically significant in the lowest tertile of the composite risk score. However, for the moderate and highest tertile of the risk score, the association between sports participation and WC (moderate risk:  $-0.29$ , 95% CI  $-0.45$  to  $-0.14$ ); high risk:  $-0.46$ , 95% CI  $-0.66$  to  $-0.25$ ) and BMI (moderate


**TABLE 2** Association between weekly sports participation and body composition by tertiles of the composite risk score.

	Low risk (n = 1269)		Moderate risk (n = 1504)		High risk (n = 1226)	
	$\beta$ (95% CI)	p-value	$\beta$ (95% CI)	p-value	$\beta$ (95% CI)	p-value
<b>WC</b>						
Unadjusted	0.06 (−0.09 to 0.21)	0.446	−0.12 (−0.28 to 0.04)	0.134	−0.33 (−0.53 to −0.14)	<b>0.001</b>
Adjusted <sup>a</sup>	−0.01 (−0.16 to 0.15)	0.947	<b>−0.29 (−0.45 to −0.14)</b>	<b>&lt;0.001</b>	<b>−0.46 (−0.66 to −0.25)</b>	<b>&lt;0.001</b>
<b>BMI</b>						
Unadjusted	0.02 (−0.04 to 0.08)	0.508	<b>−0.06 (−1.12 to −0.01)</b>	<b>0.034</b>	<b>−0.13 (−0.21 to −0.06)</b>	<b>0.001</b>
Adjusted <sup>a</sup>	0.01 (−0.04 to 0.07)	0.667	<b>−0.10 (−0.16 to −0.04)</b>	<b>&lt;0.001</b>	<b>−0.14 (−0.22 to −0.06)</b>	<b>&lt;0.001</b>

Note: Numbers in bold represent significant associations between sports participation and body composition by each tertile ( $p \leq 0.05$ ).

Abbreviations:  $\beta$ , beta-coefficients; CI, confidence interval.

<sup>a</sup>Adjusted for age, sex, ethnicity, and marital status.


**FIGURE 1** Region of significance for the association between sports participation and waist circumference depending on levels of the composite obesogenic family environment risk score.

risk:  $-0.10$ , 95% CI ( $-0.16$  to  $-0.04$ ); high risk:  $-0.14$ , 95% CI ( $-0.22$  to  $-0.06$ ), increased with higher obesogenic family risk. For example, for every one-hour increase in weekly sports participation, WC decreased approximately 0.3 cm in children from moderate obesogenic environments and 0.5 cm if the child were living in a high obesogenic family environment.

### 3.3 | Regions of significance

Figures 1 and 2 shows the point along the continuum of the composite risk score where the association between sports participation and WC or BMI transit between statistically non-significant

and significant. The distribution of participants along the moderator was 4.4% ( $-2$  SD), 27.4% ( $-1$  SD), 37.6% (mean), 25.9% (1 SD), and 4.7% (2 SD), respectively. With regards to the association between sports participation and WC, the transitional point was  $-1.15$  SD from the mean of the composite risk score, corresponding to a composite risk score of 2.4. The point where the association between sports participation and BMI transitions between non-statistically and significant was  $-0.8$  SD from the mean, equivalent to a composite risk score of 3. Thus, sports participation was only associated with WC when the composite risk score was equal to or higher than 2.4, and only significantly associated with BMI when the composite risk score equalled or was higher than 3.



**FIGURE 2** Region of significance for the association between sports participation and body mass index depending on levels of the composite obesogenic family environment risk score.

## 4 | DISCUSSION

To our knowledge, this is the first study to investigate the moderating effect of the obesogenic level of the family environment on the association between sports participation and body composition in children. We observed a statistically significant moderating effect of the obesogenic family environment on the association between sports participation and both WC and BMI, respectively. More specifically, sports participation was inversely associated with WC and BMI only among children living in a moderate-to-high risk obesogenic family environment. Furthermore, the strength of the inverse association between sports participation and body composition increased with higher obesogenic family risk score. The results suggest that sports participation may protect against obesity among those children more vulnerable or at higher risk of developing excessive weight, supporting earlier findings in the literature.<sup>21</sup>

An inverse association between sports participation and both WC and BMI have previously been demonstrated in the ENERGY project.<sup>4</sup> However, the literature has overall demonstrated ambiguous results regarding the relation between sports participation and body composition,<sup>5</sup> suggesting that this association is complex. Our main analysis showed no associations between sports participation and body composition among those children living in a low-risk obesogenic home environment. One explanation could be that a low obesogenic family home environment protects against developing overweight, independently of participating or not in sports. This would comply with previous findings indicating that the most influencing environment in regards to childhood obesity often is the home environment.<sup>7</sup> Indeed, sport programmes for children and adolescents are not necessarily designed to

prevent obesity; however, children who participate in organized sports are often more active and active at higher levels of intensities than non-participants.<sup>3</sup> Importantly, participating in sports can be beneficial for children's general physical, psychological, and social health,<sup>35,36</sup> and should be encouraged for all children.

Previous studies examining the association between an obesogenic family environment and childhood obesity have found conflicting results.<sup>8</sup> The heterogeneity of study outcomes might be a result of methodological differences in how family environmental risk factors are operationalized and measured, hampering direct comparisons between studies. Most studies have investigated only few factors or domains in the family environment related to excessive weight in children.<sup>8</sup> The present study used a composite risk score consisting of nine items primarily related to the PA, food, and media environment in the home. As also argued by others,<sup>37</sup> a composite risk score should capture the overall obesogenic family environment better than single risk factors alone. For instance, some families may exhibit both unfavourable and favourable health behaviours simultaneously, such as high levels of time in sedentary behaviours concurrently with high levels of physical activity.<sup>38</sup> A systematic review by Kininmonth et al.<sup>8</sup> investigated the association between different aspects in the home environment (related to food, PA, and media use) and child obesity, found that most studies investigated the influence of the media environment only (23 out of 62). Consequently, this could increase the risk of misclassification when defining the degree of an obesogenic family environment as well as hampering the relationship with obesity. Some of the inconsistency observed in the literature could also be attributed to the age of the included samples and individual variation in susceptibility to an obesogenic environment. For instance, the prevalence of

overweight increases with age<sup>39</sup> as well as BMI heritability.<sup>40,41</sup> This might explain why some studies in children as young as 5–6 years have not found an association between an obesogenic family environment and BMI,<sup>37</sup> compared to positive associations found in older children.<sup>42</sup> The relative contributions of genetic and social-environmental factors to body composition are controversial. Some studies indicate that obesity-related genes are more strongly associated with BMI in more obesogenic environments,<sup>19</sup> thus indicating that the observed variation in BMI heritability (31%–90%)<sup>43</sup> might be attributed to environmental characteristics. Several surveys have been utilized to assess the degree of an obesogenic family environment and how it relates to child obesity.<sup>44–46</sup> For instance the Family Nutrition and Physical Activity (FNPA) screening tool has been associated with obesity<sup>46</sup> and one-year change in BMI in children.<sup>47</sup> Both studies used the originally FNPA questionnaire based on 21 questions related to parental behaviours, rules, and support of PA, food intake, and media use in the home. Recently, the FNPA questionnaire was reworded and updated to ensure its continued relevance,<sup>48</sup> showing similar associations with BMI.<sup>49</sup> However, both the old and new version of the FNPA questionnaires also features child behaviours, like child PA levels, diet intake and sleep<sup>46</sup> and thus does not represent the obesogenic environment alone. The composite risk score used in the present study was based on fewer variables than the FNPA screening tool. However, the score was still sensitive enough to capture higher levels in WC and BMI and importantly, the variables represented aspects of the family home environment only.

There is an urgent need for effective interventions that can curb and reverse the obesity trend among children. The importance of such strategies has become even more crucial as a part of the COVID-19 pandemic recovery. Emerging studies reveals that more children has become affected by overweight and obesity during the COVID-19 lockdown,<sup>50</sup> despite recent years of stagnation in BMI trends among children in many countries.<sup>51</sup> In addition to the importance of family-based interventions to support healthy lifestyle habits for child weight management,<sup>52</sup> the present study indicates that enrolling children in sports, and especially groups of children at increased risk of overweight, should be a public health priority. Children participating in sports from an early age are not only more active, but also more prone to persist in sports and keep a lifelong interest in PA.<sup>5</sup> Thus, developing effective strategies of how to ease enrolment in sports among subgroups of children with different family backgrounds and experiences with sports, as well as reduce the high levels of drop out from sports reported with increasing age, are important from a public health perspective. In addition, we need more prospective observational and experimental studies that investigate the preventive effect of youth sports participation on obesity, especially among children at increased risk of developing overweight or obesity, for example due to their obesogenic family environment.

#### 4.1 | Strengths and limitation

Strengths of the present study include a large multinational sample from different regions across Europe, standardized data collection

protocols, and objectively measured weight, height, and waist circumference among children. We included two indicators of body composition and found the same results independent of measure used, which strengthens the conclusions of the current study.

Limitations include the cross-sectional design of the study, thus implying that causal inferences cannot be made. Further, the majority of variables were collected through questionnaires, which might have introduced bias. The obesogenic risk score is limited to nine variables and other important risk factors in an obesogenic environment might have been missed out. For instance, some studies find an association between the physical activity environment, such as sports equipment and garden space, to be related to unhealthy weight in children.<sup>53</sup> However, since this study used data from the ENERGY project, we could only use the variables that were available in the dataset. Parental BMI was, on the other hand, incorporated in the composite risk score and evaluated as a part of healthy parental role modelling. Body composition underlies a genetic influence as earlier discussed, which may have strengthened the associations with child body composition (Table S1). Another limitation is that BMI and WC are only indicators of body composition and more complex and precise methods (e.g., bioelectrical impedance, dual-energy x-ray absorptiometry, and total body water estimates) exist.<sup>54</sup> Furthermore, a summed risk score weights the variables equally, which might not need to be the case, because some variables may be more strongly related to obesity than others. Also, the obesogenic risk score was based on dichotomized variables which, together with choice of cut point, might have resulted in loss of precision. When no official recommendations were available to guide our decision of a cut-off point, we used the same cut-offs as in previous studies from the ENERGY project or based our decision on the median split. One variable, parental sugared drinks, was not correlated to either child WC or BMI (Table S1). However, since the variables were included in our composite risk score a posteriori, based on earlier findings linking the variables to increased risk of child obesity, we choose to keep this variable in the composite risk score. Adjustment for child total PA levels was not possible due to lack of information but could have revealed a more specific association between sports participation and body composition.

The generalizability of the results may have been influenced by the differences between participants included and those not included in the analysis. However, differences were small and most likely of minor importance. Finally, data was obtained in 2010 and some measures, such as screen time, lacks current popular screen use such as smart phones.

## 5 | CONCLUSION

The present study found that an obesogenic family environment modified the association between children's sports participation and body composition. Sports participation was inversely associated with WC and BMI only among children living in more obesogenic family environments, suggesting that sports participation might protect against unfavourable weight gain in this group of children. In addition to



strategies aiming to improve family environments from obesogenic to healthy, we recommend initiatives to increase interest and participation in sports from an early age, especially among subgroups of children at increased risk of excessive weight gain.

## AUTHOR CONTRIBUTIONS

Mai J.M. ChinAPaw, Yannis Manios, Dénes Molnár, Maïté Verloigne, Germàn Vicente-Rodríguez and Elling Bere contributed to the development of measurement protocols and instruments, and coordinated or supervised the data collections in the participating countries. Elling Bere, Mette Stavnsbo, and Mai J.M. ChinAPaw conceived the original idea for the manuscript and details were discussed with all authors. Mette Stavnsbo conducted the analyses with supervision from Andreas Stenling. All authors assisted in the interpretation of the results. Mette Stavnsbo drafted the manuscript, and all authors critically reviewed all drafts and have approved the submitted manuscript.

## FUNDING INFORMATION

The ENERGY-project is funded by the Seventh Framework Programme (CORDIS FP7) of the European Commission, HEALTH (FP7-HEALTH-2007-B), grant agreement no. 223254. Additional support was given from the Netherlands Organization for Health Research and Development (Grant number 50-50150-98-002). The content of this article reflects only the authors' views, and the European Community is not liable for any use that may be made of the information contained therein.

## CONFLICT OF INTEREST STATEMENT

No conflict of interest was declared.

## DATA AVAILABILITY STATEMENT

The dataset used and analysed during the current study are available from the corresponding author on reasonable request.

## ORCID

Mette Stavnsbo  <https://orcid.org/0000-0002-0920-964X>

Mai J. M. Chinapaw  <https://orcid.org/0000-0001-6259-2441>

## REFERENCES

1. Reilly JJ, Methven E, McDowell ZC, et al. Health consequences of obesity. *Arch Dis Child*. 2003;88(9):748-752. doi:10.1136/adc.88.9.748
2. Reiner M, Niermann C, Jekauc D, Woll A. Long-term health benefits of physical activity—a systematic review of longitudinal studies. *BMC Public Health*. 2013;13:813. doi:10.1186/1471-2458-13-813
3. Kokko S, Martin L, Geidne S, et al. Does sports club participation contribute to physical activity among children and adolescents? A comparison across six European countries. *Scand J Public Health*. 2019; 47(8):851-858. doi:10.1177/1403494818786110
4. Fernandez-Alvira JM, te Velde SJ, De Bourdeaudhuij I, et al. Parental education associations with children's body composition: mediation effects of energy balance-related behaviors within the ENERGY-project. *Int J Behav Nutr Phys Act*. 2013;10:80. doi:10.1186/1479-5868-10-80
5. Lee JE, Pope Z, Gao Z. The role of youth sports in promoting children's physical activity and preventing pediatric obesity: a systematic review. *Behav Med*. 2018;44(1):62-76. doi:10.1080/08964289.2016.1193462
6. Poehlman ET. A review: exercise and its influence on resting energy metabolism in man. *Med Sci Sports Exerc*. 1989;21(5):515-525.
7. Boonpleng W, Park CG, Gallo AM, Corte C, McCreary L, Bergren MD. Ecological influences of early childhood obesity: a multilevel analysis. *West J Nurs Res*. 2013;35(6):742-759. doi:10.1177/0193945913480275
8. Kininmonth AR, Smith AD, Llewellyn CH, Dye L, Lawton CL, Fildes A. The relationship between the home environment and child adiposity: a systematic review. *Int J Behav Nutr Phys Act*. 2021;18(1):4. doi:10.1186/s12966-020-01073-9
9. van Stralen MM, te Velde SJ, Singh AS, et al. European Energy balance Research to prevent excessive weight Gain among Youth (ENERGY) project: design and methodology of the ENERGY cross-sectional survey. *BMC Public Health*. 2011;11:65. doi:10.1186/1471-2458-11-65
10. Bjornara HB, Vik FN, Brug J, et al. The association of breakfast skipping and television viewing at breakfast with weight status among parents of 10–12-year-olds in eight European countries; the ENERGY (European Energy balance Research to prevent excessive weight Gain among Youth) cross-sectional study. *Public Health Nutr*. 2014; 17(4):906-914. doi:10.1017/S136898001300061X
11. Vik FN, Bjornara HB, Overby NC, et al. Associations between eating meals, watching TV while eating meals and weight status among children, ages 10-12 years in eight European countries: the ENERGY cross-sectional study. *Int J Behav Nutr Phys Act*. 2013;10:58. doi:10.1186/1479-5868-10-58
12. Cameron AJ, van Stralen MM, Brug J, et al. Television in the bedroom and increased body weight: potential explanations for their relationship among European schoolchildren. *Pediatr Obes*. 2013;8(2):130-141. doi:10.1111/j.2047-6310.2012.00094.x
13. Vik FN, Te Velde SJ, Van Lippevelde W, et al. Regular family breakfast was associated with children's overweight and parental education: results from the ENERGY cross-sectional study. *Prev Med*. 2016;91: 197-203. doi:10.1016/j.ypmed.2016.08.013
14. van Stralen MM, te Velde SJ, van Nassau F, et al. Weight status of European preschool children and associations with family demographics and energy balance-related behaviours: a pooled analysis of six European studies. *Obes Rev*. 2012;13(Suppl 1):29-41. doi:10.1111/j.1467-789X.2011.00959.x
15. Brug J, van Stralen MM, Te Velde SJ, et al. Differences in weight status and energy-balance related behaviors among schoolchildren across Europe: the ENERGY-project. *PLoS One*. 2012;7(4):e34742. doi:10.1371/journal.pone.0034742
16. Van Lippevelde W, te Velde SJ, Verloigne M, et al. Associations between home- and family-related factors and fruit juice and soft drink intake among 10- to 12-year old children. The ENERGY project. *Appetite*. 2013;61(1):59-65. doi:10.1016/j.appet.2012.10.019
17. Timperio AF, van Stralen MM, Brug J, et al. Direct and indirect associations between the family physical activity environment and sports participation among 10–12 year-old European children: testing the EnRG framework in the ENERGY project. *Int J Behav Nutr Phys Act*. 2013;10:15. doi:10.1186/1479-5868-10-15
18. te Velde SJ, ChinAPaw MJM, De Bourdeaudhuij I, et al. Parents and friends both matter: simultaneous and interactive influences of parents and friends on European schoolchildren's energy balance-related behaviours—the ENERGY cross-sectional study. *Int J Behav Nutr Phys Act*. 2014;11:82. doi:10.1186/1479-5868-11-82
19. Schrepff S, van Jaarsveld CHM, Fisher A, et al. Variation in the heritability of child body mass index by obesogenic home environment. *JAMA Pediatr*. 2018;172(12):1153-1160. doi:10.1001/jamapediatrics.2018.1508
20. Stavnsbo M, Aadland E, Anderssen SA, et al. Effects of the active smarter kids (ASK) physical activity intervention on cardiometabolic

- risk factors in children: a cluster-randomized controlled trial. *Prev Med*. 2020;130:105868. doi:10.1016/j.ypmed.2019.105868
21. Weintraub DL, Tirumalai EC, Haydel KF, Fujimoto M, Fulton JE, Robinson TN. Team sports for overweight children: the Stanford Sports to Prevent Obesity Randomized Trial (SPORT). *Arch Pediatr Adolesc Med*. 2008;162(3):232-237. doi:10.1001/archpediatrics.2007.43
  22. Nemet D, Barzilay-Teeni N, Eliakim A. Treatment of childhood obesity in obese families. *J Pediatr Endocrinol Metab*. 2008;21(5):461-467. doi:10.1515/JPEM.2008.21.5.461
  23. Chaput JP, Saunders TJ, Carson V. Interactions between sleep, movement and other non-movement behaviours in the pathogenesis of childhood obesity. *Obes Rev*. 2017;18(Suppl 1):7-14. doi:10.1111/obr.12508
  24. Singh AS, Vik FN, Chinapaw MJ, et al. Test-retest reliability and construct validity of the ENERGY-child questionnaire on energy balance-related behaviours and their potential determinants: the ENERGY-project. *Int J Behav Nutr Phys Act*. 2011;8:136. doi:10.1186/1479-5868-8-136
  25. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320(7244):1240-1243. doi:10.1136/bmj.320.7244.1240
  26. Brug J, van Stralen MM, ChinAPaw MJM, et al. Differences in weight status and energy-balance related behaviours according to ethnic background among adolescents in seven countries in Europe: the ENERGY-project. *Pediatr Obes*. 2012;7(5):399-411. doi:10.1111/j.2047-6310.2012.00067.x
  27. Velde SJT, Haraldsen E, Vik FN, et al. Associations of commuting to school and work with demographic variables and with weight status in eight European countries: the ENERGY-cross sectional study. *Prev Med*. 2017;99:305-312. doi:10.1016/j.ypmed.2017.03.005
  28. Sofi F, Cesari F, Abbate R, Gensini GF, Casini A. Adherence to Mediterranean diet and health status: meta-analysis. *BMJ*. 2008;337:a1344. doi:10.1136/bmj.a1344
  29. McNeish D, Stapleton LM, Silverman RD. On the unnecessary ubiquity of hierarchical linear modeling. *Psychol Methods*. 2017;22(1):114-140. doi:10.1037/met0000078
  30. Raftery AE. Bayesian model selection in social research. *Sociol Methodol*. 1995;25:111-163. doi:10.2307/271063
  31. Hayes AF, Matthes J. Computational procedures for probing interactions in OLS and logistic regression: SPSS and SAS implementations. *Behav Res Methods*. 2009;41(3):924-936. doi:10.3758/BRM.41.3.924
  32. Duriancik DM, Goff CR. Children of single-parent households are at a higher risk of obesity: a systematic review. *J Child Health Care*. 2019;23(3):358-369. doi:10.1177/1367493519852463
  33. Shah B, Tombeau Cost K, Fuller A, Birken CS, Anderson LN. Sex and gender differences in childhood obesity: contributing to the research agenda. *BMJ Nutr Prev Health*. 2020;3(2):387-390. doi:10.1136/bmjnp-2020-000074
  34. Sallis JF, Zakarian JM, Hovell MF, Hofstetter CR. Ethnic, socioeconomic, and sex differences in physical activity among adolescents. *J Clin Epidemiol*. 1996;49(2):125-134. doi:10.1016/0895-4356(95)00514-5
  35. Bjornara HB, Westergren T, Sejersted E, et al. Does organized sports participation in childhood and adolescence positively influence health? A review of reviews. *Prev Med Rep*. 2021;23:101425. doi:10.1016/j.pmedr.2021.101425
  36. Eime RM, Young JA, Harvey JT, Charity MJ, Payne WR. A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport. *Int J Behav Nutr Phys Act*. 2013;10:98. doi:10.1186/1479-5868-10-98
  37. Schrepft S, van Jaarsveld CH, Fisher A, Wardle J. The obesogenic quality of the home environment: associations with diet, physical activity, TV viewing, and BMI in preschool children. *PLoS One*. 2015;10(8):e0134490. doi:10.1371/journal.pone.0134490
  38. Ekelund U, Luan J, Sherar LB, et al. Moderate to vigorous physical activity and sedentary time and cardiometabolic risk factors in children and adolescents. *JAMA*. 2012;307(7):704-712. doi:10.1001/jama.2012.156
  39. Collaboration NCDRF. Height and body-mass index trajectories of school-aged children and adolescents from 1985 to 2019 in 200 countries and territories: a pooled analysis of 2181 population-based studies with 65 million participants. *Lancet*. 2020;396(10261):1511-1524. doi:10.1016/S0140-6736(20)31859-6
  40. Silventoinen K, Jelenkovic A, Sund R, et al. Genetic and environmental effects on body mass index from infancy to the onset of adulthood: an individual-based pooled analysis of 45 twin cohorts participating in the Collaborative project of Development of Anthropometrical measures in Twins (CODATwins) study. *Am J Clin Nutr*. 2016;104(2):371-379. doi:10.3945/ajcn.116.130252
  41. Silventoinen K, Rokholm B, Kaprio J, Sorensen TI. The genetic and environmental influences on childhood obesity: a systematic review of twin and adoption studies. *Int J Obes (Lond)*. 2010;34(1):29-40. doi:10.1038/ijo.2009.177
  42. MacFarlane A, Cleland V, Crawford D, Campbell K, Timperio A. Longitudinal examination of the family food environment and weight status among children. *Int J Pediatr Obes*. 2009;4(4):343-352. doi:10.3109/17477160902846211
  43. Min J, Chiu DT, Wang Y. Variation in the heritability of body mass index based on diverse twin studies: a systematic review. *Obes Rev*. 2013;14(11):871-882. doi:10.1111/obr.12065
  44. Pinard CA, Yaroch AL, Hart MH, Serrano EL, McFerren MM, Estabrooks PA. The validity and reliability of the comprehensive home environment survey (CHES). *Health Promot Pract*. 2014;15(1):109-117. doi:10.1177/1524839913477863
  45. Gattshall ML, Shoup JA, Marshall JA, Crane LA, Estabrooks PA. Validation of a survey instrument to assess home environments for physical activity and healthy eating in overweight children. *Int J Behav Nutr Phys Act*. 2008;5:3. doi:10.1186/1479-5868-5-3
  46. Ihmels MA, Welk GJ, Eisenmann JC, Nusser SM. Development and preliminary validation of a family nutrition and physical activity (FNPA) screening tool. *Int J Behav Nutr Phys Act*. 2009;6:14. doi:10.1186/1479-5868-6-14
  47. Ihmels MA, Welk GJ, Eisenmann JC, Nusser SM, Myers EF. Prediction of BMI change in young children with the family nutrition and physical activity (FNPA) screening tool. *Ann Behav Med*. 2009;38(1):60-68. doi:10.1007/s12160-009-9126-3
  48. Peyer KL, Bailey-Davis L, Welk G. Development, applications, and refinement of the family nutrition and physical activity (FNPA) child obesity prevention screening. *Health Promot Pract*. 2021;22(4):456-461. doi:10.1177/1524839920922486
  49. Peyer KL, Welk GJ. Construct validity of an obesity risk screening tool in two age groups. *Int J Environ Res Public Health*. 2017;14(4):419. doi:10.3390/ijerph14040419
  50. Salway R, Foster C, de Vocht F, et al. Accelerometer-measured physical activity and sedentary time among children and their parents in the UK before and after COVID-19 lockdowns: a natural experiment. *Int J Behav Nutr Phys Act*. 2022;19(1):51. doi:10.1186/s12966-022-01290-4
  51. Collaboration NCDRF. Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet*. 2017;390(10113):2627-2642. doi:10.1016/S0140-6736(17)32129-3
  52. Pamungkas RA, Chamroonsawasdi K. Home-based interventions to treat and prevent childhood obesity: a systematic review and meta-analysis. *Behav Sci (Basel)*. 2019;9(4):38. doi:10.3390/bs9040038
  53. Schalkwijk AAH, van der Zwaard BC, Nijpels G, Elders PJM, Platt L. The impact of greenspace and condition of the neighbourhood on child overweight. *Eur J Public Health*. 2018;28(1):88-94. doi:10.1093/eurpub/ckx037

54. Duren DL, Sherwood RJ, Czerwinski SA, et al. Body composition methods: comparisons and interpretation. *J Diabetes Sci Technol*. 2008;2(6):1139-1146. doi:[10.1177/193229680800200623](https://doi.org/10.1177/193229680800200623)

#### SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Stavnsbo M, Stenling A, Berntsen S, et al. Does an obesogenic family environment moderate the association between sports participation and body composition in children? The ENERGY project. *Pediatric Obesity*. 2023;18(7):e13031. doi:[10.1111/jjpo.13031](https://doi.org/10.1111/jjpo.13031)