

1 **Perceived exercise limitation in asthma: the role of disease severity, overweight and**
2 **physical activity in children***

3 Westergren, Thomas¹, Berntsen, Sveinung¹, Lødrup Carlsen, Karin C.^{2,3}, Mowinckel, Petter²,
4 Håland, Geir², Fegran, Liv¹, Carlsen, Kai-Håkon^{2,3}

5 ¹*Faculty of Health and Sport Sciences, University of Agder.* ²*Division of Paediatric and*
6 *Adolescent Medicine, Oslo University Hospital.* ³*Faculty of Medicine, University of Oslo.*
7 *(Norway).*
8

9 Running title: *Exercise limitation in asthma*

10 Corresponding author; Thomas Westergren

11 University of Agder, Department of Health and Nursing Science, Postbox 422, NO-4604
12 Kristiansand, NORWAY

13 Tel +47 372 33 142

14 Email thomas.westergren@uia.no

15

16 * The study was performed within ORAACLE (the Oslo Research Group of Asthma and
17 Allergy in Childhood; the Lung and Environment).

18 **Word count:** 2898

19 **Number of tables:** 2

20 **Number of figures:** 2

21

22 Westergren, T., Berntsen, S., Lødrup Carlsen, K.C., Mowinckel, P., Håland, G., Fegran, L.,
23 Carlsen, K.H.

24 **Perceived exercise limitation in asthma: the role of disease severity, overweight and**
25 **physical activity in children**

26 *Pediatr Allergy Immunol.*

27 **Abstract**

28 *Background:* Children with asthma may be less physically active than their healthy peers. We
29 aimed to investigate if perceived exercise limitation (EL) was associated with lung function or
30 bronchial hyperresponsiveness (BHR), socio-economic factors, prenatal smoking, overweight,
31 allergic disease, asthma severity or physical activity (PA).

32 *Methods:* The 302 children with asthma from the 10-year examination of the Environment and
33 Childhood Asthma birth cohort study underwent a clinical examination including perceived
34 EL (structured interview of child and parent(s)), measure of overweight (body mass index by
35 sex and age passing through 25kg/m² or above at 18 years), exercise-induced
36 bronchoconstriction (forced expiratory volume in one second (FEV₁) pre- and post-exercise),
37 methacholine bronchial challenge (severe BHR; provocative dose causing ≥20% decrease in
38 FEV₁ ≤ 1 μmol) and asthma severity score (dose of controller medication and exacerbations
39 last 12 months). Multivariate logistic regression analyses were conducted to assess
40 associations with perceived EL.

41 *Results:* In the final model explaining 30.1%, asthma severity score (OR 1.49, (1.32, 1.67))
42 and overweight (OR 2.35 (1.14, 4.82)) only were significantly associated with perceived EL.
43 Excluding asthma severity and allergic disease, severe BHR (OR 2.82 (1.38, 5.76)) or

44 maximal reduction in FEV₁ post exercise (OR 1.48 (1.10, 1.98)) and overweight (OR 2.15
45 (1.13, 4.08) and 2.53 (1.27, 5.03)) explained 9.7% and 8.4% of perceived EL, respectively.

46 *Conclusions:* Perceived EL in children with asthma was independently associated with asthma
47 severity and overweight, the latter doubling the probability of perceived EL irrespectively of
48 asthma severity, allergy status, socio-economic factors, prenatal smoking or PA.

49

50 *Keywords (MeSH):* Bronchial Hyperreactivity, Bronchial Provocation Tests, Cohort Studies,
51 Exercise-Induced Asthma, Overweight, Pulmonary Function Tests, Self-Report,
52 Socioeconomic Status

53 Thomas Westergren, University of Agder, Department of Health and Nursing Science,
54 Postboks 422, NO-4604 Kristiansand, NORWAY. Tel +47 372 33 142 Email
55 thomas.westergren@uia.no

56 ***Introduction***

57 Asthma may result in reduced physical activity (PA) and thus poor physical fitness in
58 childhood (1). Overweight children with asthma report greater limitations of PA (2).
59 Improvement in asthma control by use of long-term controller medication is shown to be
60 associated with increased PA (1). Furthermore, psychological adjustment to asthma and
61 perceived competence of PA are positively associated with increased fitness in children with
62 asthma (2, 3). Nevertheless, vigorous intensity PA may induce asthma symptoms in up to
63 90% of children with non-treated asthma and result in avoidance of PA (4).

64 The agreement between self-reported exercise-induced symptoms and objectively
65 measured exercise-induced bronchoconstriction (EIB) is reported to be poor (5-7). Panditi and
66 Silverman (7) found a weak association between children's symptom perception and EIB,
67 which was unaffected by age, gender, asthma severity, medication, habitual PA and attitudes
68 towards PA or competitiveness. Seear et al. (5) and Joyner et al. (6) revealed that only 15%
69 and 24% of children reporting exercise limitation met the criteria of EIB (decrease of $\geq 10\%$ of
70 forced expiratory volume in 1 second (FEV₁) post-exercise), respectively, explaining
71 perceived exercise limitation (EL) by low cardiorespiratory fitness rather than EIB. Johansson
72 et al. (8); however, reported a prevalence of EIB in 42% among adolescents reporting exercise
73 induced dyspnea which was significantly higher compared with controls. Nevertheless,
74 misinterpreted symptoms in symptom-based management may lead to further inactivity (5, 6),
75 and differential diagnoses may be overlooked (5, 8, 9). Moreover, overprotection by parents,
76 and children and parents' misinterpretation of regular breathlessness during vigorous intensity
77 PA may result in fear of asthma symptoms and restriction from participation in PA leading to
78 further reduced fitness (10). Additionally, household stress factors related to low
79 socioeconomic status are associated with perceived EL in children with asthma (11).

80 In the present study we aimed to investigate if perceived EL in children from a
81 prospective birth cohort study with asthma was associated with reduced lung function or
82 bronchial hyperresponsiveness (BHR), socio-economic factors, prenatal smoking, overweight,
83 allergic rhinitis (AR), atopic eczema (AE), markers of asthma severity or by PA.

84

Submitted and accepted version

85 ***Materials and Methods***

86 *Study design and subjects*

87 From the Environment and Childhood Asthma (ECA) birth cohort described elsewhere 1019
88 children attended the ten-year follow up (12). Children without suspicion of respiratory tract
89 infection for the least 4 weeks, only, were included. Two clinical examinations with measures
90 of BHR (EIB test and methacholine bronchial challenge on separate days) were performed 2–
91 7 days apart after withholding short and long acting β_2 -agonists for at least 12 and 48 h,
92 respectively, and leukotriene antagonists for 72 h. The present study comprises the 302
93 children (193 boys, 64%) with asthma out of the 1019 children examined at 10 years, who
94 were similar to the remaining 717 children without asthma with respect to height, weight,
95 household income, prenatal smoking, AR and AE.

96 *Asthma* (ever) was defined in accordance with previously reported criteria from ECA (12) by
97 at least two of the following three criteria fulfilled: (i) Dyspnoea, chest tightness and/or
98 wheezing 0-3 years and/or 4-10 years, (ii) a doctor's diagnosis of asthma and (iii) used asthma
99 medication (β_2 -agonists, sodium chromoglycate, corticosteroids, leukotriene antagonists
100 and/or aminophylline) 0-3 years and/or 4-10 years. *Allergic rhinitis* (ever) was defined with at
101 least two out of three criteria fulfilled: (i) doctor's diagnosis of rhinitis, (ii) symptoms of
102 rhinitis and (iii) treatment for eye/nose or allergy symptoms.

103 The study was approved by the Norwegian Data Inspectorate as well as the Regional
104 Committee for Medical Research Ethics in South-Eastern Norway. Written informed consent
105 to take part was obtained from guardians of the participating children.

106

107

108 *Methods*

109 *Lung function* was measured according to European standard (13) using SensorMedics Vmax
110 20c (SensorMedics Diagnostics, Yorba Linda, CA, USA), as forced expiratory flow volume
111 loops reported as FEV₁, forced vital capacity (FVC), forced expiratory flow at 25-75% of
112 FEV₁ (FEF₂₅₋₇₅) were presented as percent predicted according to Stanojevic et al. (14), in
113 addition to FEV₁/FVC.

114 *EIB tests* were conducted as 6-8 minutes treadmill run with an exercise load of 95% of
115 maximal heart rate during the last 4 minutes, following a standardized procedure (12, 15). The
116 tests were considered positive as EIB with a reduction in FEV₁ $\geq 10\%$ of baseline FEV₁ at
117 three, six, ten, fifteen or twenty minutes after running ceased. Maximal reduction (%) in FEV₁
118 ($R_{\text{Max}}\% \text{FEV}_1$) post exercise compared with baseline was calculated.

119 *Methacholine bronchial challenges* with controlled tidal ventilation were performed
120 according to American Thoracic Society (ATS) guidelines (16) by use of SPIRA® dosimeter
121 (Spira Respiratory Care Center Ltd., Hemeelinna, Finland). The starting dose was 0.05 μmol
122 and the test continued until FEV₁ was reduced by 20% or a maximum cumulative dose of 22.4
123 μmol was reached. The tests were classified as positive in two categories with a provocative
124 dose causing $\geq 20\%$ decrease in FEV₁ (PD₂₀) of $\leq 8 \mu\text{mol}$ and $\leq 1 \mu\text{mol}$ respectively; the latter
125 categorized as severe BHR.

126 *Bronchodilator response* was assessed by the highest value of FEV₁ after inhalation of
127 Salbutamol 0.5 mg per 10 kg bodyweight 20 minutes post exercise. A bronchial lability index
128 indicating of the total variability in FEV₁ (%) was calculated as the sum of the percentage
129 decrease post exercise changes in FEV₁ plus the increase after Salbutamol inhalation
130 compared to baseline (17).

131 *Structured interview* by a physician was conducted to collect data comprising
132 respiratory symptoms of the child (ISAAC questions validated in Norwegian language (18)),
133 use of medication, household income, parental education, prenatal smoking, PA and perceived
134 EL. Daily PA was assessed by a positive answer of “PA accompanied with breathlessness and
135 sweat 6-7 times each week”; hours participating in sports per week were assessed by “hours
136 participating in organized exercise each week” and; perceived EL was assessed by a positive
137 answer to the question: “present feeling that asthma restrains PA”. Self-reported daily PA and
138 hours/week in sports were correlated with objectively recorded PA at 13 year measurements,
139 described in detail elsewhere (19).

140 *Anthropometric data* were assessed measuring height with a stadiometer to nearest 0.5
141 cm and weight (Seca 709, Seca, Hamburg, Germany) to the nearest 0.1 kg wearing light
142 clothing without shoes. Overweight was defined by international cut off points for body mass
143 index by sex and age between 2 and 18 years according to Cole et al (20). Cut off points are
144 designed to pass through 25kg/m² at age 18 years.

145 *An asthma severity score* ranging from 0-9 was constructed based upon steps
146 suggested by Taylor et al. (21). This score included the reported asthma controller medication
147 (inhaled corticosteroids (ICS) and leukotriene antagonists and/or β_2 -agonists) in addition to
148 exacerbations reported during the last 12 months (classified as 0, 1-3, and >3). Description of
149 the asthma severity score is given in table 1.

150 *Statistical analysis*

151 Chi-square tests were conducted to compare frequencies of categorical variables between
152 children with and without asthma and between the groups with and without perceived EL.
153 Continuous normally distributed variables are presented as mean with standard deviation
154 (SD). Independent t-tests were used to analyze differences between groups. Skewed variables

155 are presented as median with interquartile range (IQR), and Mann-Whitney Wilcoxon tests
156 were preferred for calculating differences in not normally distributed data.

157 Variables with a significance level of ≤ 0.20 from bivariate analyses were considered
158 for multivariate logistic regression analysis. Stepwise multivariate logistic regression analysis
159 according to Hosmer et al. (22) was conducted, removing the least significant variable until
160 only significant values remained. Results from logistic regression models are presented as
161 Odds ratio (OR) with 95% confidence interval (CI).

162 Multivariate analysis was conducted in five separate models including one measure of
163 BHR in each: EIB; $R_{\text{Max}}\%FEV_1$; $PD_{20} \leq 1 \mu\text{mol}$; $PD_{20} \leq 8 \mu\text{mol}$ and; Bronchial lability index.
164 To avoid multi-collinearity, each model included only one lung function variable with the
165 lowest p -value (FEV_1/FVC). Analysis of each model was repeated twice; first excluding
166 asthma severity markers from the total models as severity was significantly associated with all
167 five BHR variables. Secondly, allergic disease was additionally excluded from the reduced
168 model due to significant associations with both PD_{20} variables and Bronchial lability index.
169 All significant independent variables in each final model as well as gender were checked for
170 interaction terms with perceived EL.

171 Statistical significance level was set to 5%. Nagelkerke R^2 was reported as explained
172 variance from logistic regression models. Statistical analyses were performed with Statistical
173 Package for Social Sciences Version 22.0 (SPSS, Chicago, IL, USA).

174

175 **Results**

176 Fifty eight (20%) children with asthma reported EL. As shown in table 2, these children had
177 significantly larger $R_{Max}\%FEV_1$ post exercise as well as significantly more often severe BHR,
178 more often a mother who smoked during pregnancy, overweight and comorbidity of both AR
179 and AE than children without perceived EL. The asthma severity score was significantly
180 higher and FEV_1/FVC lower in children with compared to without perceived EL. A non-
181 significant ($p=0.07$) tendency to higher rate of positive EIB was found among children with
182 (35%) compared to without (24%) perceived EL (Table 2). Groups did not differ significantly
183 with regard to participation in sports or daily PA. Positive EIB-test, $PD_{20} \leq 1$, $PD_{20} \leq 8 \mu\text{mol}$,
184 Bronchial lability index $\geq 10\%$ or $FEV_1/FVC \leq 80\%$ individually or combined ranged from
185 29-67% in children with perceived EL.

186 In the final model, including gender, BHR, FEV_1/FVC , low household income,
187 prenatal smoking, overweight, allergic disease, asthma severity score and hours/week in
188 sports; asthma severity score (1.49 (1.32, 1.67)) and overweight (2.35 (1.14, 4.84)) only were
189 independently associated with perceived EL (Fig. 1a). In the model 30.1% of the variance was
190 explained. In the reduced model excluding asthma severity score 15.5% of the variance in EL
191 was explained (Fig. 1b). Children more likely to report EL were those whose mother smoked
192 during pregnancy (1.95 (1.02, 3.80)), overweight children (2.54 (1.30, 4.95)), children with
193 AR only (3.03 (1.06, 8.69)), and children with comorbidity of AR and AE (5.53 (2.49,
194 12.31)).

195 BHR was significantly associated with perceived EL only when excluding both
196 asthma severity score and allergic disease from analysis. Children with severe BHR were 2.82
197 (1.38, 5.76) times more likely (Fig. 2a), and children with 10 percent increased $R_{Max}\%FEV_1$

198 were 1.48 (1.10, 1.98) times more likely (Fig. 2b) to report EL, respectively. These models
199 explained 9.7% and 8.4% of the variance in EL.

200 Overweight children were more than twice likely to report EL with significant
201 associations to EL in all multivariate analysis (OR between 2.15 (1.13, 4.08) and 2.54 (1.30,
202 4.95)). This association was not significantly influenced by BHR, lung function, low
203 household income, maternal prenatal smoking, allergic diseases, asthma severity score or
204 hours per week in sports. Overweight did not significantly influence the associations between
205 perceived EL and asthma severity score, comorbidity of AR and AE or BHR. There were no
206 significant interactions between independent variables and perceived EL in the reported
207 models.

208

Submitted and accepted version

209 **Discussion**

210 Exercise limitation was reported in 20% of children with asthma. In the final model
211 explaining 30.1%, asthma severity score and overweight only were significantly associated
212 with perceived EL. Excluding asthma severity and allergic disease, 9.7% and 8.4% of
213 perceived EL were explained by significant associations to overweight and severe BHR or
214 $R_{\max}\%FEV_1$ post exercise, respectively. Overweight children were more than twice likely to
215 report EL irrespectively of any other included factor.

216 The 20% of children reporting EL is lower than the general activity limitation reported
217 in 52% of children with current asthma (23), although the frequency of EIB (35%) among
218 children reporting EL in the present study was comparable to previously reported range
219 between 8-42% in studies comparing self- reported symptoms with EIB test(5, 6, 8, 9, 24).
220 Nevertheless, 67 out of 264 (25%) had a positive EIB compared to previously reported 51-
221 55% (25, 26) including also children with current asthma and ICS treatment. The variations in
222 rate of reported EL or exercise-induced symptoms and associations between EIB and
223 perceived exercise-induced symptoms (5-8) may be related to study populations, asthma
224 definitions and control. Children with asthma ever in the present birth cohort study are likely
225 to have less severe asthma (27) than the current asthma patients in the study by Yeatts et al.
226 (23). Also, different assessments may affect the rate of EL or EIB, with our children
227 responding to a question whether they experienced that their asthma restricted their physical
228 activity, compared to limited activities because of wheezing ≥ 1 times per month.

229 Despite no significant association between a positive EIB and perceived EL, the
230 associations between $R_{\max}\%FEV_1$ as well as severe BHR and EL suggest that perceived EL
231 reflects BHR. This is supported by a report by Sanchez-Garcia et al. (24) suggesting that the
232 direct methacholine challenge and the indirect mannitol tests have high sensitivities to detect

233 BHR in steroid naive children complaining of one or more symptoms after exercise. Sanchez-
234 Garcia et al. (24) reported a detection rate of BHR in 96.7% with a methacholine test,
235 increasing to 100% when combined with mannitol test (24). In contrast, Anderson et al. (28)
236 reported a sensitivity of 59% and 56%, and a specificity of 65% and 69% to identify
237 objectively measured EIB by mannitol and methacholine, respectively (28). The American
238 Thoracic Society guidelines recommend mannitol test or hyperosmolar aerosols of 4.5%
239 saline or eucapnic voluntary hyperpnoea of dry air as surrogates of exercise test (29), although
240 these were not performed in the present study. Nevertheless, neither methacholine bronchial
241 challenge, nor EIB-test individually or combined with $FEV_1/FVC \leq 80\%$ or Bronchial lability
242 index $\leq 10\%$ confirmed EIB in more than 67% of children reporting EL. This may be related
243 to the anti-inflammatory BHR reducing effect of ICS (16), used by many of our study subjects
244 compared to the steroid naive children in the study by Sanchez-Garcia et al. (24). The
245 associations between perceived EL and asthma severity and allergic disease may additionally
246 reflect the impact of uncontrolled asthma.

247 In the models excluding severity and allergic disease, explained variation of perceived
248 EL was 8.4% to 9.7%. Asthma severity score and overweight adjusted for gender; however,
249 statistically explained 30.1% of the variance in reported EL, and in the model excluding
250 severity; 15.5% of the variance were explained without contribution from objective
251 measurements. Both asthma severity score and allergic disease, which are clinically accessible
252 without objective measures, were hence advantageous to objective measures in explained
253 variation of reported EL. Moreover, contrary to Panditi and Silverman (7) who found no
254 association of severity and perception of exercise induced symptoms, our findings confirmed
255 that severity assessed objectively by BHR or qualitatively by a severity score was related to
256 perceived EL.

257 We found no significant associations between reported PA and perceived EL, in line
258 with previous studies (2, 9), possibly indicating an absence of felt limitation due to inactivity
259 or participation in PA despite perceived EL. Children and their parents may avoid symptoms
260 through less PA, through overprotection (10) and/or beliefs that asthma limits the possibility
261 for PA (30). On the other hand, we previously showed in the ECA study by objective
262 recordings of PA at 13 years of age that children were rather active (19, 27), indicating that
263 children in the present study might have been rather active despite 80% reporting PA less
264 frequently than daily.

265 The robust association between perceived EL and overweight was in contrast to the
266 report by Joyner et al. (6) who found no association between BMI and self-reported EL.
267 Pianosi and Davis (2); however, reported that overweight children with asthma perceived
268 greater limitations of PA. Causal relationship explaining why overweight may induce
269 limitations is complex as asthma symptoms may induce perceived EL, reduced PA level and
270 thus development of overweight (1). We were not able to verify whether low
271 cardiorespiratory fitness, as suggested in previous reports (5, 6, 9), may explain perceived EL
272 in children with asthma even without BHR. However, similar fitness and PA level in
273 overweight and normal weight children with asthma are previously reported (2). We hence
274 interpret overweight as an independent perceived barrier of PA which may be labeled to
275 asthma by children, irrespectively of reduced lung function, BHR, low household income,
276 prenatal smoking, allergic disease, asthma severity score or hours/week in sports.

277 *Strengths and limitations*

278 The main strengths of the present study were the nested case-control design, and the
279 assessment of perceived EL related to objective assessment of BHR and lung function through
280 validated and standardized procedures. Lack of objective measures of PA as well as fitness,

281 and flow volume loops during exercise indicating expiratory flow limitations are considered
282 as a limitation, in addition to lack of parents' and children's reports of attitudes and beliefs
283 about PA. Also, we asked if the child experienced exercise limitation or not, and did not
284 include any information on grading of limitation. We were therefore unable to identify if
285 marked EL correlated better with objective measures than did limited degree of EL. It should
286 be underlined that results from a population based study will differ from a study based on
287 patients referred to specialized service as the study by Vahlquist and Pedersen (1), which
288 included patients with more severe asthma whereas the present population based study will
289 differ as far as severity and consequences of the disease is concerned.

290 **Conclusion**

291 Perceived EL in children with asthma was independently associated with asthma severity and
292 overweight, the latter doubling the probability of perceived EL irrespectively of asthma
293 severity, allergy status, socio-economic factors, prenatal smoking or PA.

294 **Acknowledgements**

295 The present study was supported by grants from the Eastern Norway Regional Health
296 Authority with supplemental funds from the Norwegian Asthma and Allergy Association and
297 the Norwegian Nurses Organisation.

298 **Declaration of interest**

299 There is no competing or conflict of interests.

300

301 **References**

- 302 1. Vahlkvist S, Inman MD, Pedersen S. Effect of asthma treatment on fitness, daily
303 activity and body composition in children with asthma. *Allergy*. 2010;65:1464-1471.
- 304 2. Pianosi PT, Davis HS. Determinants of physical fitness in children with asthma.
305 *Pediatrics*. 2004;113:e225-229.
- 306 3. Strunk RC, Mrazek DA, Fukuhara JT, Masterson J, Ludwick SK, LaBrecque JF.
307 Cardiovascular fitness in children with asthma correlates with psychologic functioning of the
308 child. *Pediatrics*. 1989;84:460-464.
- 309 4. Milgrom H, Taussig LM. Keeping children with exercise-induced asthma active.
310 *Pediatrics*. 1999;104:e38.
- 311 5. Seear M, Wensley D, West N. How accurate is the diagnosis of exercise induced
312 asthma among Vancouver schoolchildren? *Arch Dis Child*. 2004;90:898-902.
- 313 6. Joyner BL, Fiorino EK, Matta-Arroyo E, Needleman JP. Cardiopulmonary exercise
314 testing in children and adolescents with asthma who report symptoms of exercise-induced
315 bronchoconstriction. *J Asthma*. 2006;43:675-678.
- 316 7. Panditi S, Silverman M. Perception of exercise induced asthma by children and their
317 parents. *Arch Dis Child*. 2003;88:807-811.
- 318 8. Johansson H, Norlander K, Berglund L, et al. Prevalence of exercise-induced
319 bronchoconstriction and exercise-induced laryngeal obstruction in a general adolescent
320 population. *Thorax*. 2015;70:57-63.
- 321 9. Abu-Hasan M, Tannous B, Weinberger M. Exercise-induced dyspnea in children and
322 adolescents: if not asthma then what? *Ann Allergy Asthma Immunol*. 2005;94:366-371.
- 323 10. Williams B, Hoskins G, Pow J, Neville R, Mukhopadhyay S, Coyle J. Low exercise
324 among children with asthma: a culture of over protection? A qualitative study of experiences
325 and beliefs. *Br J Gen Pract*. 2010;60:319-326.

- 326 11. Quinn K, Kaufman JS, Siddiqi A, Yeatts KB. Stress and the city: housing stressors are
327 associated with respiratory health among low socioeconomic status Chicago children. *J Urban*
328 *Health*. 2010;87:688-702.
- 329 12. Carlsen KCL, Håland G, Devulapalli CS, et al. Asthma in every fifth child in Oslo,
330 Norway: a 10-year follow up of a birth cohort study. *Allergy*. 2006;61:454-460.
- 331 13. Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir*
332 *J*. 2005;26:319-338.
- 333 14. Stanojevic S, Wade A, Stocks J, et al. Reference ranges for spirometry across all ages:
334 a new approach. *Am J Respir Crit Care Med*. 2008;177:253-260.
- 335 15. Carlsen KH, Engh G, Mørk M. Exercise-induced bronchoconstriction depends on
336 exercise load. *Respir Med*. 2000;94:750-755.
- 337 16. Crapo RO, Casaburi R, Coates AL, et al. Guidelines for methacholine and exercise
338 challenge testing-1999. This official statement of the American Thoracic Society was adopted
339 by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med*. 1999;161:309-329.
- 340 17. Remes ST, Pekkanen J, Remes K, Salonen RO, Korppi M. In search of childhood
341 asthma: questionnaire, tests of bronchial hyperresponsiveness, and clinical evaluation.
342 *Thorax*. 2002;57:120-126.
- 343 18. Selnes A, Bolle R, Holt J, Lund E. Cumulative incidence of asthma and allergy in
344 north-Norwegian schoolchildren in 1985 and 1995. *Pediatr Allergy Immunol*. 2002;13:58-63.
- 345 19. Westergren T, Carlsen KCL, Carlsen KH, Mowinckel P, Fegran L, Berntsen S. Basic
346 self-reports as an indicator of physical activity. *Eur Respir J*. 2016;48.
- 347 20. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for
348 child overweight and obesity worldwide: international survey. *BMJ*. 2000;320:1240-1243.
- 349 21. Taylor DR, Bateman ED, Boulet LP, et al. A new perspective on concepts of asthma
350 severity and control. *Eur Respir J*. 2008;32:545-554.

- 351 22. Hosmer DW, Lemeshow S, Sturdivant RX. *Applied Logistic Regression*. 3rd ed. ed:
352 United States: John Wiley & Sons Inc; 2013.
- 353 23. Yeatts K, Shy C, Sotir M, Music S, Herget C. Health consequences for children with
354 undiagnosed asthma-like symptoms. *Arch Pediatr Adolesc Med*. 2003;157:540-544.
- 355 24. Sánchez-García S, Rodríguez del Río P, Escudero C, García-Fernández C, Ibáñez MD.
356 Exercise-induced bronchospasm diagnosis in children. Utility of combined lung function
357 tests. *Pediatr Allergy Immunol*. 2015;26:73-79.
- 358 25. Anthracopoulos MB, Fouzas S, Papadopoulos M, et al. Physical activity and exercise-
359 induced bronchoconstriction in Greek schoolchildren. *Pediatr Pulmonol*. 2012;47:1080-1087.
- 360 26. Waalkens HJ, van Essen-Zandvliet EE, Gerritsen J, Duiverman EJ, Kerrebijn KF,
361 Knol K. The effect of an inhaled corticosteroid (budesonide) on exercise-induced asthma in
362 children. Dutch CNSLD Study Group. *Eur Respir J*. 1993;6:652-656.
- 363 27. Berntsen S, Carlsen KCL, Anderssen SA, et al. Norwegian adolescents with asthma
364 are physical active and fit. *Allergy*. 2009;64:421-426.
- 365 28. Anderson SD, Charlton B, Weiler JM, Nichols S, Spector SL, Pearlman DS.
366 Comparison of mannitol and methacholine to predict exercise-induced bronchoconstriction
367 and a clinical diagnosis of asthma. *Respir Res*. 2009;10:4.
- 368 29. Parsons JP, Hallstrand TS, Mastrorarde JG, et al. An official American Thoracic
369 Society clinical practice guideline: exercise-induced bronchoconstriction. *Am J Respir Crit*
370 *Care Med*. 2013;187:1016-1027.
- 371 30. Williams B, Powell A, Hoskins G, Neville R. Exploring and explaining low
372 participation in physical activity among children and young people with asthma: a review.
373 *BMC Fam Pract*. 2008;9:1-11.
- 374

375 **Table 1** Description of asthma severity score based on the use and dose of inhaled corticosteroids, use of
376 leukotriene antagonists/ β_2 -agonists and number of exacerbations last 12 months.

	Dose of ICS (μg)			Use of LKTR and/or β_2 -agonists		Number of exacerbations last 12 months			Total possible score
	0-99	100-399	≥ 400	yes	no	0	1-3	>3	
Score	0	2	4	1	0	0	2	4	0-9

377

378 Abbreviations: ICS; inhaled corticosteroids, LKTR; leukotriene antagonist

379

Submitted and accepted version

380 **Table 2** Descriptive characteristics of children with asthma reporting or not reporting exercise limitation (EL).
 381 Data are given as count (%) including n=294 unless otherwise stated.

	Perceived EL (n=58)	Not perceived EL (n=236)	p-value ^a
Boys ^b	35 (60)	152 (64)	0.57
EIB ($\geq 10\%$ fall in FEV ₁) ⁽ⁿ⁼²⁶⁴⁾	18 (35)	49 (23)	0.07
R _{Max} %FEV ₁ , median (IQR) ⁽ⁿ⁼²⁶⁴⁾	7 (10)	6 (6)	0.02
Bronchial lability index (%), median (IQR) ⁽ⁿ⁼²⁵³⁾	10 (12)	9 (8)	0.09
Severe BHR (PD ₂₀ $\leq 1\mu\text{mol}$) ⁽ⁿ⁼²⁹¹⁾	17 (29)	33 (14)	0.01
BHR (PD ₂₀ ≤ 8) ⁽ⁿ⁼²⁹¹⁾	31 (53)	101 (43)	0.17
FEV ₁ (% of predicted), mean (SD) ⁽ⁿ⁼²⁹⁰⁾	94 (11)	96 (9)	0.12
FVC (% of predicted), mean (SD) ⁽ⁿ⁼²⁹⁰⁾	99 (10)	98 (9)	0.91
FEV ₁ /FVC, mean (SD) ⁽ⁿ⁼²⁹⁰⁾	83 (7)	84 (6)	0.04
FEF ₂₅₋₇₅ (% of predicted), mean (SD) ⁽ⁿ⁼²⁹⁰⁾	81 (22)	86 (19)	0.14
Low household income (< 350.000NOK /year) ^{c (n=290)}	16 (28)	40 (17)	0.07
Low parental education (no education beyond 13 years of schooling)	22 (38)	88 (37)	0.93
Prenatal smoking	20 (35)	50 (21)	0.03
Overweight	21 (36)	46 (20)	0.01
Allergic rhinitis only	7 (12)	23 (10)	0.60
Atopic eczema only	14 (24)	60 (25)	0.84
Allergic rhinitis + atopic eczema	24 (41)	43 (18)	<0.01
Asthma severity score, median (IQR)	5 (5)	0 (3)	<0.01
Use of ICS last 12 months	37 (64)	46 (20)	<0.01
Hours / week in sports, median (IQR)	2 (3)	2 (2)	0.16
Daily physical activity	13 (22)	41 (18)	0.39

382

383 ^a statistical differences between groups in bold. Analysis conducted: categorical variables; Chi-square tests,
 384 continuous normally distributed variables; independent t-tests, skewed variables; Mann-Whitney Wilcoxon tests.
 385 Statistical significant differences at 5% level are given in bold.

386 ^b with reference to girls

387 ^c corresponding to ≈ 43000 €

388 Abbreviations: n; numbers, EIB; exercise induced bronchoconstriction, FEV₁; forced expiratory volume in 1
 389 second, R_{Max}%FEV₁; maximal reduction in FEV₁ post exercise (%), IQR; interquartile range, BHR; bronchial
 390 hyper responsiveness, SD; standard deviation, FVC; forced vital capacity, FEF₂₅₋₇₅; forced expiratory flow at 25-
 391 75% of FEV₁, NOK; Norwegian Kroner, ICS; inhaled corticosteroids.


392

Figure 1 Radar plots visualizing Odds ratio (OR) of factors associated with perceived exercise limitation (EL) in children with asthma showing (A) a total model with explained variance of 30.1% in EL including a composite Asthma severity score (use of asthma medication and asthma exacerbations) and (B) a model excluding the Asthma severity score with explained variance of 15.5% in EL. Numbers from 0 to 2.5 (A) and 0 to 6 (B) illustrate the OR for reporting EL. Models were adjusted for gender. The least significant variables were removed stepwise according to Hosmer et al (22) until only significant variables remained.


Submitted and accepted version

Fig. 1.

(A)



(B)




Abbreviations: BHR; bronchial hyper responsiveness, FEV₁; forced expiratory volume in 1 second, FVC; forced vital capacity

Figure 2 Radar plots visualizing Odds ratio (OR) of factors associated with perceived exercise limitation (EL) in children with asthma using Severe BHR (B) and $R_{Max}\%FEV_1$ (A) to assess BHR in separate reduced multivariate logistic regression models excluding asthma severity and allergic disease from analysis. Model including Severe BHR (A) explained 9.7%, and model including $R_{Max}\%FEV_1$ (B) explained 8.4% of the variance in EL, respectively. Numbers from 0 to 3 illustrate the Odds Ratio for reporting EL. Models were adjusted for gender. The least significant variables were removed stepwise according to Hosmer et al (22) until only significant variables remained.


Submitted and accepted version

Fig. 2.

(A)



(B)



Abbreviations: FEV₁; forced expiratory volume in 1 second, FVC; forced vital capacity, R_{Max}%FEV₁; maximal reduction in FEV₁ post exercise (%), BHR; bronchial hyper responsiveness.